For Research Use Only

AGPAT2 Recombinant antibody

Catalog Number:83349-3-RR



Purification Method:

Basic Information

Catalog Number: GenBank Accession Number:

83349-3-RR BC019292 Protein A purification

Size:GeneID (NCBI):CloneNo.:100ul , Concentration: 1000 ug/ml by10555240153C2

 Nanodrop;
 UNIPROT ID:
 Recommended Dilutions:

 Source:
 015120
 WB 1:2000-1:10000

 Rabbit
 Full Name:
 IHC 1:200-1:800

Isotype:1-acylglycerol-3-phosphate O-IgGacyltransferase 2 (lysophosphatidic

Immunogen Catalog Number: acid acyltransferase, beta)

AG34761 Calculated MW:

31 kDa Observed MW: 27 kDa

Applications

Tested Applications:

WB, IHC, FC (Intra), ELISA

Species Specificity:

human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

WB: HeLa cells, Jurkat cells, THP-1 cells

IHC: human ovary cancer tissue,

Background Information

1-acyl-sn-glycerol-3-phosphate acyltransferase beta (AGPAT2) belongs to a family of enzymes catalyzing the sn-2 acylation of the glycerol-3-phosphate backbone. AGPAT2 is highly expressed in adipose tissues, liver, and skeletal muscle. AGPAT2 is the only AGPAT isoform whose loss-of-function mutations cause a severe form of human congenital generalized lipodystrophy (PMID: 34824276). Human and mouse AGPAT2 have a calculated molecular mass of 31 kDa, and an alternatively spliced form of AGPAT2 mRNA, encoding a protein of 246 rather than 278 amino acids, is also found in human (PMID: 19336658). Western detected AGPAT2 at an apparent molecular mass of 27 kDa.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 83349-3-RR (AGPAT2 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human ovary cancer tissue slide using 83349-3-RR (AGPAT2 antibody) at dilution of 1:400 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



1x10^6 A549 cells were intracellularly stained with 0.25 ug AGPAT2 Recombinant antibody (83349-3-RR, Clone:240153C2) and APC-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (red), or 0.25 ug Isotype Control (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



Biolayer interferometry (BLI) kinetic assays of 83349-3-RR against Human AGPAT2 were performed. The affinity constant is 2.72 nM.