

For Research Use Only

XRN2 Recombinant antibody

Catalog Number: 83378-1-RR



Basic Information

| | | |
|--|--|---|
| Catalog Number: 83378-1-RR | GenBank Accession Number: BC006417 | Purification Method: Protein A purification |
| Size: 100ul , Concentration: 1000 ug/ml by Nanodrop; | GeneID (NCBI): 22803 | CloneNo.: 240065G4 |
| Source: Rabbit | UNIPROT ID: Q9H0D6 | Recommended Dilutions: WB 1:5000-1:50000 |
| Isotype: IgG | Full Name: 5'-3' exoribonuclease 2 | |
| Immunogen Catalog Number: AG30781 | Calculated MW: 104 kDa | |
| | Observed MW: 109 kDa | |

Applications

| | |
|--|---|
| Tested Applications: WB, FC (Intra), ELISA | Positive Controls: WB : MCF-7 cells, Jurkat cells, HeLa cells |
| Species Specificity: human | |

Background Information

XRN2 is one exonuclease that degrades the Pol II associated product of poly(A) site cleavage, which is crucial for Pol II termination. During transcription termination, XRN2 cleavages at the polyadenylation site liberates a 5' fragment which is subsequently processed to form the mature mRNA and a 3' fragment which remains attached to the elongating polymerase. The processive degradation of this 3' fragment by this protein may promote termination of transcription.

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 83378-1-RR (XRN2 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



1x10⁶ HeLa cells were intracellularly stained with 0.25 ug XRN2 Recombinant antibody (83378-1-RR, Clone:240065G4) and APC-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L)(red), or 0.25 ug Isotype Control (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).



Biolayer interferometry (BLI) kinetic assays of 83378-1-RR against Human XRN2 were performed. The affinity constant is 1.49 nM.