For Research Use Only

PMS1 Recombinant antibody

Catalog Number:83456-6-RR




Basic Information	Catalog Number: 83456-6-RR	GenBank Accession Number BC096331	er: Purification Method: Protein A purfication	
	Size: 100ul , Concentration: 1000 ug/ml by Nanodrop; Source: Rabbit Isotype: IgG	GenelD (NCBI): 5378	CloneNo.: 240408D9	
		UNIPROT ID: P54277 Full Name:	Recommended Dilutions: WB 1:500-1:2000 IHC 1:50-1:500	
		PMS1 postmeiotic segregation increased 1 (S. cerevisiae)	tion IF/ICC 1:500-1:2000	
	Immunogen Catalog Number: AG28132	Observed MW: 106 kDa		
Applications	Tested Applications:	Pos	itive Controls:	
	WB, IHC, IF/ICC, FC (Intra), ELISA Species Specificity:		WB : HeLa cells, HepG2 cells, HEK-293T cells, K-562 cells, Jurkat cells, MOLT-4 cells, HL-60 cells	
	Note-IHC: suggested antigen retrieval with		: human ovary cancer tissue,	
			CC : HeLa cells, Caco-2 cells	
Background Information	insertions and deletions as a consequ MLH1 could form a heterodimer MutLf complex recognizes the DNA mispair: RFC, Exol were recruited, inducing Ex (PMID:24981171). Mutations in this ge	ence of DNA polymerase er 8 that belongs to MMREs (PM 5 firstly, and then the Mlh1-F onuclease 1 (Exo1)-depende ene cause hereditary nonpol ported that PMS1 is a widely	tes (MMREs) which is to eliminate the mismatch or rors at DNA synthesis in eukaryotes. PMS1 and ID:25619773). In the MMR system, the MutS Pms1 and other associated proteins such as PCNA, ent and -independent MMR pathways yposis colorectal cancer(HNPCC), which is also expressed protein and located in the nucleus.	
Background Information	insertions and deletions as a consequ MLH1 could form a heterodimer MutLf complex recognizes the DNA mispair: RFC, Exol were recruited, inducing Ex (PMID:24981171). Mutations in this ge known as Lynch syndrome. It was repu	ence of DNA polymerase er 8 that belongs to MMREs (PM 5 firstly, and then the Mlh1-F onuclease 1 (Exo1)-depende ene cause hereditary nonpol orted that PMS1 is a widely orms of PMS1 protein.	rors at DNA synthesis in eukaryotes. PMS1 and ID:25619773). In the MMR system, the MutS Pms1 and other associated proteins such as PCNA, ent and -independent MMR pathways yposis colorectal cancer(HNPCC), which is also	


For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)E: proteintech@ptglab.comW: ptglab.comW: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 83456-6-RR (PMS1 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human ovary cancer tissue slide using 83456-6-RR (PMS1 antibody) at dilution of 1:200 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using PMS1 antibody (83456-6-RR, Clone: 240408D9) at dilution of 1:1000 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-phalloidin (red).





1x10^6 HeLa cells were intracellularly stained with 0.25 ug PMS1 Recombinant antibody (83456-6-RR, Clone:240408D9) and APC-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L)(red), or 0.25 ug Isotype Control (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).

Biolayer interferometry (BLI) kinetic assays of 83456-6-RR against Human PMS1 were performed. The affinity constant is 0.367 nM.