For Research Use Only

HSPA2 Recombinant antibody

Catalog Number:83466-1-RR Featured Product




Basic Information	Catalog Number: 83466-1-RR	GenBank Accession Number: BC036107	Purification Method: Protein A purification					
	Size: 100ul , Concentration: 1000 ug/ml by Nanodrop; Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG3539	GenelD (NCBI):	CloneNo.: 240421A2 Recommended Dilutions: WB 1:5000-1:50000 IHC 1:250-1:1000					
		3306						
		UNIPROT ID: P54652 Full Name:						
				heat shock 70kDa protein 2 Calculated MW: 639 aa, 70 kDa Observed MW: 70 kDa				
		Applications	Tested Applications: WB, IHC, ELISA Species Specificity: human, mouse				Positive Controls: WB : HeLa cells, A431 cells, A549 cells, HaCaT cells, NIH/3T3 cells, human kidney tissue IHC : mouse testis tissue,	
				NIH/313 0				
			Note-IHC: suggested antigen ro TE buffer pH 9.0; (*) Alternativ retrieval may be performed w buffer pH 6.0	vely, antigen				
	Human HSPA2 is a member of the HSPA (HSP70) family of heat-shock proteins, encoded by the gene originally described as testis-specific. Recently, it has been reported that HSPA2 can be also expressed in human somatic tissues in a cell-type specific manner. HSPA2 is a cytosol/nuclear protein able to translocate between cytoplasm and nucleus. At physiological temperature, the HSPA2 was localized primarily in cytoplasm whereas, during heat shock, localization shifted to nucleus and nucleoli (PMID: 18452162). HSPA2 is involved in sperm maturity, function and fertility. Aberrant expression of HSPA2 in testes has been connected with male infertility. Recently, HSPA2 has attracted increased interest due to its possible involvement in carcinogenesis of non-testicular tissues. This antibody well recognized the endogenous HSPA2 protein in testis and multiple cell lines. (21373891)							
Background Information	described as testis-specific. Recently, tissues in a cell-type specific manner and nucleus. At physiological temper shock, localization shifted to nucleus and fertility. Aberrant expression of H attracted increased interest due to its	it has been reported that HSPA2 c HSPA2 is a cytosol/nuclear prote ature, the HSPA2 was localized pri and nucleoli (PMID: 18452162). H ISPA2 in testes has been connecte possible involvement in carcinog	an be also expressed in human somatic in able to translocate between cytoplasm marily in cytoplasm whereas, during heat SPA2 is involved in sperm maturity, functior d with male infertility. Recently, HSPA2 has enesis of non-testicular tissues. This					
Background Information	described as testis-specific. Recently, tissues in a cell-type specific manner, and nucleus. At physiological temper shock, localization shifted to nucleus and fertility. Aberrant expression of H attracted increased interest due to its antibody well recognized the endoge Storage: Store at -20°C. Stable for one year after Storage Buffer: PBS with 0.02% sodium azide and 50°	it has been reported that HSPA2 c. HSPA2 is a cytosol/nuclear prote ature, the HSPA2 was localized pri and nucleoli (PMID: 18452162). H ISPA2 in testes has been connecte possible involvement in carcinog nous HSPA2 protein in testis and n er shipment. % glycerol pH 7.3.	an be also expressed in human somatic in able to translocate between cytoplasm marily in cytoplasm whereas, during heat SPA2 is involved in sperm maturity, functior d with male infertility. Recently, HSPA2 has enesis of non-testicular tissues. This					
	described as testis-specific. Recently, tissues in a cell-type specific manner, and nucleus. At physiological temper shock, localization shifted to nucleus and fertility. Aberrant expression of H attracted increased interest due to its antibody well recognized the endoge Storage: Storage: Storage Buffer:	it has been reported that HSPA2 c. HSPA2 is a cytosol/nuclear prote ature, the HSPA2 was localized pri and nucleoli (PMID: 18452162). H ISPA2 in testes has been connecte possible involvement in carcinog nous HSPA2 protein in testis and n er shipment. % glycerol pH 7.3.	an be also expressed in human somatic in able to translocate between cytoplasm marily in cytoplasm whereas, during heat SPA2 is involved in sperm maturity, function d with male infertility. Recently, HSPA2 has enesis of non-testicular tissues. This					


For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data




WB result of HSPA2 antibody (83466-1-RR; 1:1000; incubated at room temperature for 1.5 hours) with sh-Control and sh-HSPA2 transfected HeLa cells.




Various lysates were subjected to SDS PAGE followed by western blot with 83466-1-RR (HSPA2 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded mouse testis tissue slide using 83466-1-RR (HSPA2 antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse testis tissue slide using 83466-1-RR (HSPA2 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Biolayer interferometry (BLL) kinetic assays of 83466-1-RR against Human HSPA2 were performed. The affinity constant is 1.86 nM.