For Research Use Only

Neurogranin Recombinant antibody

Catalog Number:83593-5-RR



Purification Method:

Protein A purfication

Recommended Dilutions:

WB 1:2000-1:10000

IHC 1:250-1:1000

CloneNo.:

240526B9

Basic Information

Catalog Number: GenBank Accession Number:

83593-5-RR BC002835 GeneID (NCBI): Size:

Nanodrop: **UNIPROT ID:** Q92686 Rabbit Full Name:

100ul, Concentration: 1000 ug/ml by 4900

Isotype: neurogranin (protein kinase C substrate, RC3)

IgG Immunogen Catalog Number: Calculated MW: AG0676 7.6 kDa

Observed MW: 15-17 kDa

Applications

Tested Applications: WB, IHC, FC (Intra), ELISA

Species Specificity:

human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

WB: mouse brain tissue, rat brain tissue

IHC: mouse brain tissue,

Background Information

Neurogranin (formerly designated p17, also known as Ng, RC3 and BICKS) belongs to the neurogranin family. It is a neuron-specific substrate for protein kinase C (PKC). It is a postsynaptic protein that is highly enriched in brain, with restricted expression in the cortex, striatum, hippocampus, thalamus, hypothalamus and olfactory bulb nuclei. Neurogranin binds calmodulin at low levels of calcium, thereby regulating calmodulin-dependent nitric oxide synthase. Neurogranin acts as a "third messenger" substrate of protein kinase C-mediated molecular cascades during synaptic development and remodeling. It binds to calmodulin in the absence of calcium.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com W: ptglab.com

Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 83593-5-RR (Neurogranin antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 83593-5-RR (Neurogranin antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 83593-5-RR (Neurogranin antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



1x10^6 A549 cells were intracellularly stained with 0.25 ug Neurogranin Recombinant antibody (83593-5-RR, Clone:240526B9) and APC-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (red), or 0.25 ug Isotype Control (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



Biolayer interferometry (BLI) kinetic assays of 83593-5-RR against Human NRGN were performed. The affinity constant is 0.304 nM.