

For Research Use Only

PHKG2 Recombinant antibody

Catalog Number: 83663-2-RR



Basic Information

Catalog Number: 83663-2-RR	GenBank Accession Number: BC002541	Purification Method: Protein A purification
Size: 100ul , Concentration: 1000 ug/ml by Nanodrop;	GeneID (NCBI): 5261	CloneNo.: 240643E1
Source: Rabbit	UNIPROT ID: P15735	Recommended Dilutions: WB 1:2000-1:10000 IHC 1:200-1:800 IF/ICC 1:50-1:500
Isotype: IgG	Full Name: phosphorylase kinase, gamma 2 (testis)	
Immunogen Catalog Number: AG6829	Calculated MW: 46 kDa	
	Observed MW: 46 kDa	

Applications

Tested Applications: WB, IHC, IF/ICC, FC (Intra), ELISA	Positive Controls: WB : HepG2 cells, MCF-7 cells, mouse liver tissue IHC : mouse liver tissue, IF/ICC : HEK-293 cells,
Species Specificity: human, mouse	
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	

Background Information

PHKG2 (phosphorylase kinase catalytic subunit gamma 2), it is expected to be located in the cytoplasm, which is enriched in the testis. Catalytic subunit of the phosphorylase b kinase (PHK), which mediates the neural and hormonal regulation of glycogen breakdown (glycogenolysis) by phosphorylating and thereby activating glycogen phosphorylase. It may regulates glycogeneolysis in the testis. Phosphorylase kinase is a polymer of 16 subunits, four each of alpha, beta, gamma and delta. The gamma subunit also includes the skeletal muscle and hepatic isoforms, and the hepatic isoform is encoded by this gene. The gamma subunits contain the active site of the enzyme, whereas the alpha and beta subunits have regulatory functions controlled by phosphorylation. The delta subunit mediates the dependence of the enzyme on calcium concentration. The molecular weight of PHKG2 is 46 kDa.

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA




For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 83663-2-RR (PHKG2 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.




Immunofluorescent analysis of (-20°C Ethanol) fixed HEK-293 cells using PHKG2 antibody (83663-2-RR, Clone: 240643E1) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2).



1x10⁶ HEK-293T cells were intracellularly stained with 0.25 ug PHKG2 Recombinant antibody (83663-2-RR, Clone:240643E1) and APC-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L)(red), or 0.25 ug Isotype Control (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



1x10⁶ HepG2 cells were intracellularly stained with 0.25 ug PHKG2 Recombinant antibody (83663-2-RR, Clone:240643E1) and APC-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L)(red), or 0.25 ug Isotype Control (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



Biolayer interferometry (BLI) kinetic assays of 83663-2-RR against Human PHKG2 were performed. The affinity constant is 1.90 nM.