

For Research Use Only

CDCA7L Recombinant antibody

Catalog Number: 83682-4-RR



Basic Information

Catalog Number: 83682-4-RR	GenBank Accession Number: BC009352	Purification Method: Protein A purification
Size: 100ul , Concentration: 1000 ug/ml by Nanodrop;	GeneID (NCBI): 55536	CloneNo.: 240639A3
Source: Rabbit	UNIPROT ID: Q96GN5	Recommended Dilutions: WB 1:500-1:2000
Isotype: IgG	Full Name: cell division cycle associated 7-like	
Immunogen Catalog Number: AG10944	Calculated MW: 454 aa, 52 kDa	
	Observed MW: 52 kDa	

Applications

Tested Applications: WB, FC (Intra), ELISA	Positive Controls: WB : A549 cells, HEK-293T cells, HeLa cells, K-562 cells, MCF-7 cells
Species Specificity: human	

Background Information

Cell division cycle-associated 7-like protein (CDCA7L) is also known as HR1 and JPO2 and belongs to the JPO protein family. CDCA7L is identified as a target gene of c-Myc and is frequently dysregulated in multiple cancers. CDCA7L can complement the c-Myc transformation-defective mutant W135E and potentiate Myc-mediated transformation (PMID: 24141559, PMID: 37140023). CDCA7L translocates from cytoplasm to nucleus under dexamethasone induction. CDCA7L is widely expressed. CDCA7L has 4 isoforms produced by alternative splicing (refer to UniProt).

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 83682-4-RR (CDCA7L antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



1x10⁶ A431 cells were intracellularly stained with 0.25 ug CDCA7L Recombinant antibody (83682-4-RR, Clone:240639A3) and APC-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (red), or 0.25 ug Isotype Control (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).



Biolayer interferometry (BLI) kinetic assays of 83682-4-RR against Human CDCA7L were performed. The affinity constant is 7.75 nM.