For Research Use Only

## CD34 Recombinant antibody

Catalog Number:83713-2-RR





| Basic Information               | Catalog Number:<br>83713-2-RR   | GenBank Accession Number:<br>NM_133654              | Purification Method:<br>Protein A purification                  |  |
|---------------------------------|---|---|---|--|
|                                 | Size:<br>100ul , Concentration: 1000 µg/ml by<br>Nanodrop;<br>Source:<br>Rabbit<br>Isotype:<br>IgG  | GenelD (NCBI):<br>12490                             | CloneNo.:<br>240659C7   |  |
|                                 |   | UNIPROT ID:<br>Q64314<br>Full Name:<br>CD34 antigen | Recommended Dilutions:<br>WB 1:5000-1:50000<br>IHC 1:500-1:2000 |  |
|                                 |   |   | IF/ICC 1:125-1:500  |  |
|                                 |   | Calculated MW:<br>41 kDa                            |   |  |
|                                 |   | Observed MW:<br>105 kDa                             |   |  |
| Applications                    | Tested Applications:<br>WB, IHC, IF/ICC, ELISA  | Positi  | ive Controls:   |  |
|                                 | Species Specificity:  |   | C2C12 cells,<br>mouse testis tissue,                            |  |
|                                 | Note-IHC: suggested antigen retrieval with<br>TE buffer pH 9.0; (*) Alternatively, antigen<br>retrieval may be performed with citrate<br>buffer pH 6.0  |   | C : NIH/3T3 cells,  |  |
| Background Information          | CD34 is a 105- to 120-kDa glycophosphoprotein expressed on the majority of hematopoietic stem/progenitor cells,<br>bone marrow stromal cells, capillary endothelial cells, embryonic fibroblasts, and some nerve tissue. CD34 is a<br>commonly used marker for identifying human hematopoietic stem/progenitor cells and mediates cell adhesion and<br>lymphocyte homing by binding L-selectin and E-selectin ligands. CD34 is also one of the best negative selection<br>markers for characterizing and/or isolating human MSCs from bone marrow and other sources. Along with other<br>positive selection markers (such as CD29, CD44, CD90, CD105 and CD166), negative selection markers (such as<br>CD34 and CD45) are used for MSC identification. |   |   |  |
| Storage                         | Storage:<br>Store at -20°C. Stable for one year aft<br>Storage Buffer:<br>PBS with 0.02% sodium azide and 50<br>Aliquoting is unnecessary for -20°C s   | % glycerol pH 7.3.                                  |   |  |
| *** 20ul sizes contain 0.1% BSA | Arequiring to uninecessary for -20 C 3  | wide.   |   |  |

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free<br/>in USA), or 1(312) 455-8498 (outside USA)E: proteintech@ptglab.comW: ptglab.comW: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data






Various lysates were subjected to SDS PAGE followed by western blot with 83713-2-RR (CD34 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.

Immunofluorescent analysis of (4% PFA) fixed NIH/3T3 cells using CD34 antibody (83713-2-RR, Clone: 240659C7) at dilution of 1:250 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2).

Immunohistochemical analysis of paraffinembedded mouse testis tissue slide using 83713-2-RR (CD34 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Biolayer interferometry (BLI) kinetic assays of 83713-2-RR against Mouse CD34 were performed. The affinity constant is 1.09 nM.