For Research Use Only

NEU2 Recombinant antibody, PBS Only

Catalog Number:83934-1-PBS



Purification Method:

Protein A purfication

CloneNo.:

240685B3

Basic Information

Catalog Number: GenBank Accession Number:

83934-1-PBS

GeneID (NCBI): Size:

100ug, Concentration: 1 mg/ml by 4759 Nanodrop; **UNIPROT ID:** Source: Q9Y3R4

Isotype: sialidase 2 (cytosolic sialidase)

IgG Calculated MW: Immunogen Catalog Number: 380 aa, 42 kDa AG17386 Observed MW:

42 kDa

Full Name:

Applications

Tested Applications: WB, IHC, ELISA

Rabbit

Species Specificity: human, mouse

Background Information

NEU2, also named as Sialidase 2, neuraminidase 2, catalyzes the hydrolytic cleavage of the terminal sialic acid (Nacetylneuraminic acid, Neu5Ac) of a glycan moiety in the catabolism of glycolipids, glycoproteins and oligosaccharides. NEU2 is localized in the cytosol and plasma membrane and takes part in myoblast and neuronal differentiation. It is almost undetectable under normal physiological conditions, and was found to be involved into apoptotic events in pancreatic ductal adenocarcinoma, one of the most dangerous cancers due to poor diagnosis and prognosis. (PMID: 30439363)


Storage

Storage:

Store at -80°C. Storage Buffer: PBS only

in USA), or 1(312) 455-8498 (outside USA)

Selected Validation Data




mouse skeletal muscle tissue were subjected to SDS PAGE followed by western blot with 83934-1-RR (NEU2 antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 83934-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded mouse skeletal muscle tissue slide using 83934-1-RR (NEU2 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 83934-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded mouse skeletal muscle tissue slide using 83934-1-RR (NEU2 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 83934-1-PBS in a different storage buffer formulation.



Biolayer interferometry (BLI) kinetic assays of 83934-1-RR against Human NEU2 were performed. The affinity constant is 2.76 nM.