For Research Use Only

CCAR1 Recombinant antibody

Catalog Number:83936-4-RR





Basic Information	Catalog Number: 83936-4-RR	GenBank Accession Number: BC132725 GeneID (NCBI): 55749 UNIPROT ID: Q8IX12 Full Name: cell division cycle and apoptosis regulator 1 Calculated MW: 1150 aa, 133 kDa		Purification Method: Protein A purfication CloneNo.: 240980C12 Recommended Dilutions: WB 1:5000-1:50000 IHC 1:200-1:800 IF/ICC 1:125-1:500				
	100ul , Concentration: 1000 ug/ml by Nanodrop; Source: Rabbit Isotype: IgG							
						Observed MW: 150 kDa		
					Applications	Tested Applications:		Positive Controls:
	WB, IHC, IF/ICC, ELISA Species Specificity:	Jurkat cells		ells, HEK-293 cells, HeLa cells, HepG2 cells, n colon tissue,				
human								
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0		IF/ICC : A431	cells,					
Background Information	Cell cycle and apoptosis regulator 1 (CCAR1) was originally identified from a functional screening and implicated in retinoid-induced cell cycle arrest and apoptosis. It has been shown that, in breast cancer cells, CCAR1 regulates apoptosis induced by various agents, including chemotherapeutics adriamycin, etoposide, and EGFR inhibitor. It is also involved in mediating anchorage-independent growth of colon carcinoma cells. In addition, CCAR1 plays the role as a co-activator of nuclear receptors to recruit mediator complex. It was found to associate with components of the Mediator and p160 coactivator complexes and is recruited to endogenous NR target genes. CCAR1 is also a binding partner of b-catenin and assists b-catenin in transcriptional activation of Wnt target genes.							
	Storage:	er shinment						
Storage	Store at -20°C. Stable for one year after Storage Buffer: PBS with 0.02% sodium azide and 50° Aliquoting is unnecessary for -20°C st	% glycerol						

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com


This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





Various lysates were subjected to SDS PAGE followed by western blot with 83936-4-RR (CCAR1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. Interpret to the subject of the



Immunohistochemical analysis of paraffinembedded human colon tissue slide using 83936-4-RR (CCAR1 antibody) at dilution of 1:400 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).

Immunohistochemical analysis of paraffinembedded human colon tissue slide using 83936-4-RR (CCAR1 antibody) at dilution of 1:400 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).