For Research Use Only

CXXC5 Recombinant antibody

Catalog Number:83940-7-RR




Basic Information	Catalog Number: 83940-7-RR	GenBank Accession Number: BC017439	Purification Method: Protein A purfication
	100ul , Concentration: 1000 µg/ml by Nanodrop; Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG9733		CloneNo.: 241034A6 Recommended Dilutions: WB 1:5000-1:50000
		UNIPROT ID: Q7LFL8 Full Name: CXXC finger 5 Calculated MW: 322aa,33 kDa; 227aa,24 kDa Observed MW: 33-40 kDa	
Applications	Tested Applications: WB, ELISA Species Specificity: human	Positive Controls: WB : MCF-7 cells, K-562 cells, mouse kidney tissue	
Background Information	CXXC5 (CXXC-type zinc finger protein 5) is also named as HSPC 195, RINF and TCCCIA00297.CXXC5 is a member of the CXXC-type zinc-finger protein family. CXXC5 is a short protein (322 amino acids in length) that does not have any catalytic domain, but is able to bind to DNA and act as a transcription factor and epigenetic factor through protein-protein interactions. In addition, CXXC5 is capable of regulating various signal transduction processes, including the TGF-β, Wnt and ATM-p53 pathways, thereby acting as a novel and crucial signaling coordinator. CXXC5 plays an important role in embryonic development and adult tissue homeostasis by regulating cell proliferation, differentiation and apoptosis (PMID: 30479059). CXXC5 has two isoforms of 33 kDa and 24 kDa (PMID: 25805812).		
Storage	Storage: Store at -20°C. Stable for one year aft Storage Buffer: PBS with 0.02% sodium azide and 50 Aliguoting is unnecessary for -20°C s	% glycerol pH 7.3.	
*** 20ul sizes contain 0.1% BSA	,	U	

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)E: proteintech@ptglab.comW: ptglab.comW: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 83940-7-RR (CXXC5 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.