

For Research Use Only

# Coagulation factor II Recombinant antibody

Catalog Number: 83957-4-RR



## Basic Information

Catalog Number:	83957-4-RR	GenBank Accession Number:	NM_010168.3	Purification Method:	Protein A purification
Size:	100ul , Concentration: 1000 ug/ml by Nanodrop;	GeneID (NCBI):	14061	CloneNo.:	241013D9
Source:	Rabbit	UNIPROT ID:	P19221	Recommended Dilutions:	WB: 1:2000-1:10000
Isotype:	IgG	Full Name:	coagulation factor II		
Immunogen Catalog Number:	EG0943	Calculated MW:	70 kDa		
		Observed MW:	70-80 kDa		

## Applications

Tested Applications:	WB, ELISA	Positive Controls:	
Species Specificity:	mouse, rat	WB :	mouse heart tissue, rat plasma

## Background Information

Normal blood coagulation is a complex process, involving a cascade of activation of different plasma proteins, ultimately resulting in the formation of a clot, called fibrin. Coagulation Factor II (F2), also known as prothrombin or factor II, is one of the components of this chain of plasma proteins involved in blood coagulation. Prothrombin is the precursor of thrombin, which is essential in the processes of hemostasis and thrombosis. This antibody specifically recognizes mouse Coagulation Factor II.

## Storage


**Storage:**  
Store at -20°C. Stable for one year after shipment.  
**Storage Buffer:**  
PBS with 0.02% sodium azide and 50% glycerol, pH7.3  
Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:  
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)  
E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 83957-4-RR (Coagulation factor II antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Biolayer interferometry (BLI) kinetic assays of 83957-4-RR against Mouse Coagulation factor II were performed. The affinity constant is 16.4 nM.