

For Research Use Only

NMI Recombinant antibody

Catalog Number: 83986-2-RR

Featured Product



Basic Information

Catalog Number:

83986-2-RR

Size:

100ul , Concentration: 1000 µg/ml by Nanodrop;

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG34759

GenBank Accession Number:

BC001268

GeneID (NCBI):

9111

UNIPROT ID:

Q13287

Full Name:

N-myc (and STAT) interactor

Calculated MW:

35 kDa

Observed MW:

35 kDa

Purification Method:

Protein A purification

CloneNo.:

241135A9

Recommended Dilutions:

WB 1:5000-1:50000

IF/ICC 1:200-1:800

Applications

Tested Applications:

WB, IF/ICC, FC (Intra), ELISA

Species Specificity:

human

Positive Controls:

WB : HeLa cells, K-562 cells, PC-3 cells, HepG2 cells

IF/ICC : HepG2 cells,

Background Information

N-Myc and STAT Interactor protein (NMI) is an interferon inducible protein that interacts with NMYC and CMYC (members of the oncogene Myc family), and other transcription factors containing a Zip, HLH, or HLH-Zip motif. It is widely involved in the process of tumor growth, progression, and metastasis.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




WB result of NMI antibody (83986-2-RR; 1:2000; incubated at room temperature for 1.5 hours) with sh-Control and sh-NMI transfected HeLa cells.




Various lysates were subjected to SDS PAGE followed by western blot with 83986-2-RR (NMI antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using NMI antibody (83986-2-RR, Clone: 241135A9) at dilution of 1:400 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2).



1x10⁶ HepG2 cells were intracellularly stained with 0.25 ug NMI Recombinant antibody (83986-2-RR, Clone:241135A9) and APC-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L)(red), or 0.25 ug Isotype Control (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



Biolayer interferometry (BLI) kinetic assays of 83986-2-RR against Human NMI were performed. The affinity constant is 1.98 nM.