

For Research Use Only

NMI Recombinant antibody, PBS Only

Catalog Number: 83986-5-PBS



Basic Information

Catalog Number:

83986-5-PBS

Size:

100ug , Concentration: 1 mg/ml by Nanodrop;

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG34759

GenBank Accession Number:

BC001268

GeneID (NCBI):

9111

UNIPROT ID:

Q13287

Full Name:

N-myc (and STAT) interactor

Calculated MW:

35 kDa

Observed MW:

37 kDa

Purification Method:

Protein A purification

CloneNo.:

241135H2

Applications

Tested Applications:

WB, IF/ICC, FC (Intra), ELISA

Species Specificity:

human

Background Information

N-Myc and STAT Interactor protein (NMI) is an interferon inducible protein that interacts with NMYC and CMYC (members of the oncogene Myc family), and other transcription factors containing a Zip, HLH, or HLH-Zip motif. It is widely involved in the process of tumor growth, progression, and metastasis.

Storage

Storage:

Store at -80°C.

Storage Buffer:

PBS Only

For technical support and original validation data for this product please contact:


T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com


W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 83986-5-RR (NMI antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 83986-5-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (-20°C Ethanol) fixed HeLa cells using NMI antibody (83986-5-RR, Clone: 241135H2) at dilution of 1:400 and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2). This data was developed using the same antibody clone with 83986-5-PBS in a different storage buffer formulation.



1x10⁶ HepG2 cells were intracellularly stained with 0.25 ug NMI Recombinant antibody (83986-5-RR, Clone:241135H2) and APC-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L)(red), or 0.25 ug Isotype Control (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same antibody clone with 83986-5-PBS in a different storage buffer formulation.



Biolayer interferometry (BLI) kinetic assays of 83986-5-RR against Human NMI were performed. The affinity constant is 5.17 nM.