For Research Use Only

ZNF148 Recombinant antibody

Catalog Number:84338-7-RR



Basic Information

Catalog Number: GenBank Accession Number:

84338-7-RR BC035591 GeneID (NCBI): Size:

100ul , Concentration: 1000 $\mu g/ml$ by 7707

Nanodrop: **UNIPROT ID:** Q9UQR1 Rabbit

Full Name: Isotype: zinc finger protein 148

IgG Observed MW: Immunogen Catalog Number: 103 kDa

AG35742

human

Applications

Tested Applications:

WB, ELISA Species Specificity: Positive Controls:

WB: MCF-7 cells, T-47D cells, HCT 116 cells, HeLa cells, MDA-MB-231 cells, HepG2 cells, NIH/3T3 cells

Purification Method:

Protein A purfication

Recommended Dilutions:

WB 1:2000-1:10000

CloneNo.:

241632D8

Background Information

The Kruppel-like zinc finger protein 148 (ZNF148), also known as β enolase repressor factor 1 (BERF1, ZBP-89, and BFCOL1), encoded by this gene is a member of the Kruppel family of zinc finger DNA binding proteins. ZNF 148 participates in the regulation of cell proliferation and death, and is expressed at lower levels in mammalian somatic cells, and highly expressed in tumor cells. Previous studies show that there was a significant correlation between ZNF148 expression and the invasion and metastasis of colorectal tumors. ZNF148 has two alternative splicing isoforms, which are ZNF148FL and ZNF148AN. ZNF148FL contains a complete 794 amino acids, ZNF148AN lacks the amino-terminal 129 amino acids, and the mechanisms involved in the generation of these two splicing isoforms are not yet clear (PMID: 30463804). Thus, ZNF148 can be observed to have two molecular weights near 103

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


*** 20ul sizes contain 0.1% BSA

in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 84338-7-RR (ZNF 148 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.