

For Research Use Only

Acetyl-Histone H2B (Lys120) Recombinant antibody, PBS Only

Catalog Number: 84551-1-PBS



Basic Information

Catalog Number:

84551-1-PBS

Size:

100ug , Concentration: 1 mg/ml by
Nanodrop;

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

BC005827

GeneID (NCBI):

8349

UNIPROT ID:

Q16778

Full Name:

histone cluster 2, H2be

Calculated MW:

14 kDa

Observed MW:

25 kDa

Purification Method:

Protein A purification

CloneNo.:

241556E7

Applications

Tested Applications:

WB, IF/ICC, Dot Blot, Indirect ELISA

Species Specificity:

human, mouse

Background Information

Histones are nuclear proteins that are classified into five major protein groups: histones H2A, H2B, H3, and H4 are known as the core histones. Post-translationally modified H2B proteins can modulate the nucleosome/chromatin structure or DNA accessibility to affect the transcriptional pathways linked to embryonic development and cell differentiation. Monoubiquitination of histone H2B has emerged as an important chromatin modification with roles not only in transcription but also in cell differentiation, DNA repair or mRNA processing(PMID: 25027370).

Storage

Storage:

Store at -80°C.

Storage Buffer:

PBS Only


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Dot blot analysis was used to confirm the specificity of Acetyl-Histone H2B (Lys120) antibody. Peptides were spotted onto NC and probed with antibody at 1 µg/ml. The amount of peptide (µg/mL) spotted is indicated next to each row. This data was developed using the same antibody clone with 84551-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed Trichostatin A treated HeLa cells using Acetyl-Histone H2B (Lys120) antibody (84551-1-RR, Clone: 241556E7) at dilution of 1:1000 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-Phalloidin (red). This data was developed using the same antibody clone with 84551-1-PBS in a different storage buffer formulation.



Trichostatin A treated NIH/3T3 cells and untreated cells were subjected to SDS PAGE followed by western blot with 84551-1-RR (Acetyl-Histone H2B (Lys120) antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 84551-1-PBS in a different storage buffer formulation.