

For Research Use Only

OSMR Recombinant antibody, PBS Only

Catalog Number: 84555-2-PBS



Basic Information

Catalog Number:

84555-2-PBS

Size:

100ug , Concentration: 1 mg/ml by Nanodrop;

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG1426

GenBank Accession Number:

BC010943

GeneID (NCBI):

9180

UNIPROT ID:

Q99650

Full Name:

oncostatin M receptor

Calculated MW:

110 kDa

Observed MW:

~180 kDa

Purification Method:

Protein A purification

CloneNo.:

241865C8

Applications

Tested Applications:

WB, IHC, FC (Intra), Indirect ELISA

Species Specificity:

human, mouse

Background Information

Oncostatin M (OSM) is a member of interleukin-6 family cytokines that modulates proliferation and differentiation in numerous cells (PMID: 22982441). OSM signals via a receptor complex comprising gp130 and the OSMR (also called OSMR β) (PMID: 8999038). OSMR is a single-pass membrane protein that belongs to the type I cytokine receptor family. OSMR mRNA is expressed at relatively high levels in all neural cells and fibroblast, epithelial, and various tumor cell lines (PMID: 8999038). It has also been shown that OSMR associates with IL31RA to form the heterodimeric IL31 receptor (PMID: 15184896). OSMR has a calculated molecular weight of 110 kDa, with an apparent molecular weight of 150-180 kDa (PMID: 8999038; 17028186; 39266541; 34644107). A 70 kDa soluble form can be detected by western blot (PMID: 17028186).

Storage

Storage:

Store at -80°C.

Storage Buffer:

PBS Only


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 84555-2-RR (OSMR antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 84555-2-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffin-embedded mouse cerebellum tissue slide using 84555-2-RR (OSMR antibody) at dilution of 1:400 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 84555-2-PBS in a different storage buffer formulation.



1x10⁶ HeLa cells were intracellularly stained with 0.25 ug OSMR Recombinant antibody (84555-2-RR, Clone:241865C8) and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2)(red), or 0.25 ug Rabbit IgG Isotype Control RecAb (98136-1-RR, Clone: 240953C9) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same antibody clone with 84555-2-PBS in a

