For Research Use Only

OSMR Recombinant antibody

Catalog Number:84555-2-RR



Basic Information

Catalog Number: GenBank Accession Number:

84555-2-RR BC010943 GeneID (NCBI): Size:

100ul , Concentration: 1000 $\mu g/ml$ by 9180 Nanodrop: **UNIPROT ID:** Q99650 Rabbit Full Name:

Isotype: oncostatin M receptor IgG Calculated MW: Immunogen Catalog Number: 110 kDa AG1426 Observed MW:

~180 kDa

Applications

Tested Applications:

WB, IHC, FC (Intra), ELISA

Species Specificity:

human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Purification Method: Protein A purfication

CloneNo.: 241865C8

Positive Controls:

Recommended Dilutions:

WB 1:500-1:2000 IHC 1:200-1:800

WB: HeLa cells, Raji cells, A375 cells, HEK-293 cells,

HepG2 cells, SH-SY5Y cells, NIH/3T3 cells

IHC: mouse cerebellum tissue,

Background Information

Oncostatin M (OSM) is a member of interleukin-6 family cytokines that modulates proliferation and differentiation in numerous cells (PMID: 22982441). OSM signals via a receptor complex comprising gp130 and the OSMR (also called OSMRβ) (PMID: 8999038). OSMR is a single-pass membrane protein that belongs to the type I cytokine $receptor\ family.\ OSMR\ mRNA\ is\ expressed\ at\ relatively\ high\ levels\ in\ all\ neural\ cells\ and\ fibroblast,\ epithelial,\ and\ neural\ cells\ and\ fibroblast,\ epithelial\ and\ expressed\ at\ relatively\ high\ levels\ in\ all\ neural\ cells\ and\ fibroblast,\ epithelial\ and\ expressed\ at\ relatively\ high\ levels\ in\ all\ neural\ cells\ and\ fibroblast,\ epithelial\ and\ expressed\ at\ relatively\ high\ levels\ in\ all\ neural\ cells\ and\ fibroblast,\ epithelial\ and\ expressed\ at\ relatively\ high\ levels\ in\ all\ neural\ expressed\ at\ relatively\ high\ levels\ high\ high\ levels\ high\ levels\ high\ hig$ various tumor cell lines (PMID: 8999038). It has also been shown that OSMR associates with IL31RA to form the heterodimeric IL31 receptor (PMID: 15184896). OMSR has a calculated molecular weight of 110 kDa, with an apparent molecular weight of 150-180 kDa (PMID: 8999038; 17028186; 39266541; 34644107). A 70 kDa soluble form can be detected by western blot (PMID: 17028186).

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 84555-2-RR (OSMR antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded mouse cerebellum tissue slide using 84555-2-RR (OSMR antibody) at dilution of 1:400 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



1x10^6 HeLa cells were intracellularly stained with 0.25 ug OSMR Recombinant antibody (84555-2-RR, Clone:241865C8) and Coralite®488-Conjugated Goat Anti-Rabbit 1gG(H+L) (5A00013-2)(red), or 0.25 ug Rabbit 1gG Isotype Control Recold (98136-1-RR, Clone: 240953C9) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).