

For Research Use Only

# CEBP Alpha/CEBPA Recombinant antibody

Catalog Number: 84668-4-RR



## Basic Information

<b>Catalog Number:</b> 84668-4-RR	<b>GenBank Accession Number:</b> NM_004364	<b>Purification Method:</b> Protein A purification
<b>Size:</b> 100ul , Concentration: 1000 ug/ml by Nanodrop;	<b>GeneID (NCBI):</b> 1050	<b>CloneNo.:</b> 242101C9
<b>Source:</b> Rabbit	<b>UNIPROT ID:</b> P49715	<b>Recommended Dilutions:</b> IF/ICC 1:1000-1:4000
<b>Isotype:</b> IgG	<b>Full Name:</b> CCAAT/enhancer binding protein (C/EBP), alpha	
<b>Immunogen Catalog Number:</b> AG29947	<b>Calculated MW:</b> 38 kDa	
	<b>Observed MW:</b> 40-45 kDa	

## Applications

<b>Tested Applications:</b> IF/ICC, FC (Intra), ELISA	<b>Positive Controls:</b> IF/ICC : HepG2 cells, HeLa cells
<b>Species Specificity:</b> human	

## Background Information

CEBPA and its isoforms play important roles in lineage determination and gene activation in a variety of cell types by activating transcription from lineage-specific promoters. CEBPA is a DNA-binding protein that recognizes two different motifs: the CCAAT homology common to many promoters and the enhanced core homology common to many enhancers. In hematopoiesis, C/EBPα is a key factor in driving the development of myeloid cells interacting with a variety of factors, including c-Myc, PU.1, and microRNAs. It can also form heterodimers with the related proteins CEBP-beta and CEBP-gamma. The encoded protein has been shown to bind to the promoter and modulate the expression of the gene encoding leptin which plays an important role in body weight homeostasis. CEBPA can interact with CDK2 and CDK4, thereby inhibiting these kinases and causing growth arrest in cultured cells. Several pathways have been implicated as the means by which CEBPA mediates cell cycle arrest and proliferation, including p21, cyclin-dependent kinases and the E2F complex via c-Myc.

## Storage


**Storage:**  
Store at -20°C. Stable for one year after shipment.  
**Storage Buffer:**  
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.  
Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:  
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)  
E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data




Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using CEBP Alpha/CEBPA antibody (84668-4-RR, Clone: 242101C9) at dilution of 1:2000 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-Phalloidin (red).



1x10<sup>6</sup> HeLa cells were intracellularly stained with 0.25 ug CEBP Alpha/CEBPA Recombinant antibody (84668-4-RR, Clone:242101C9) and APC-Conjugated Goat Anti-Rabbit IgG(H+L)(red), or 0.25 ug Rabbit IgG Isotype Control Recombinant Antibody (98136-1-RR, Clone: 240953C9) (blue). Cells were fixed and permeabilized with True-Nuclear Transcription Factor Buffer Set.



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using CEBP Alpha/CEBPA antibody (84668-4-RR, Clone: 242101C9) at dilution of 1:2000 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-Phalloidin (red).



Biolayer interferometry (BLI) kinetic assays of 84668-4-RR against Human CEBP Alpha/CEBPA were performed. The affinity constant is 1.85 nM.