For Research Use Only

TMIGD2/CD28H Recombinant antibody

Catalog Number:84727-5-RR



Purification Method:

Basic Information

Catalog Number: GenBank Accession Number:

84727-5-RR BC015655 Protein A purfication

 Size:
 GeneID (NCBI):
 CloneNo.:

 100ul , Concentration: 1000 µg/ml by 126259
 241807C12

Nanodrop; UNIPROT ID: Recommended Dilutions: Source: Q96BF3 WB 1:5000-1:50000

Rabbit Full Name:

Isotype: transmembrane and immunoglobulin

IgG domain containing 2

Calculated MW: 282 aa, 31 kDa Observed MW: 48 kDa

Applications

Tested Applications:

WB : Jurkat cells, pig thymus tissue, MOLT-4 cells, HeLa

cells

Positive Controls:

Species Specificity: human, pig

Background Information

Transmembrane and immunoglobulin domain-containing protein 2 (TMIGD2, also known as CD28H and IGPR1), plays a role in cell-cell interaction, migration, and angiogenesis. Through interaction with HHLA2, it costimulates T-cells in the context of TCR-mediated activation and enhances T-cell proliferation and cytokine production via an AKT-dependent signaling cascade (PMID: 22419821; 23784006). TMIGD2 has been associated with several diseases, particularly in the context of cancer. It is aberiarly expressed by human AML cells and can be used to identify and enrich functional LSCs (PMID: 38167704). The predicted molecular weight of the TMIGD2 protein is 31 kDa. The higher molecular weight of TMIGD2 could be associated with its potential glycosylation at its extracellular region (PMID: 22419821).

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 84727-5-RR (TMIGD2/CD28H antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



pig thymus tissue were subjected to SDS PAGE followed by western blot with 84727-5-RR (TMIGD2/CD28H antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Biolayer interferometry (BLL) kinetic assays of 84727-5-RR against Human TMIGD2/CD28H were performed. The affinity constant is 1.23 nM.