For Research Use Only

## Mouse Endoglin/CD105 Recombinant antibody, PBŠ Only

Catalog Number:84807-4-PBS



**Purification Method:** 

Protein A purfication

CloneNo.:

242305G3

**Basic Information** 

Catalog Number: 84807-4-PBS

GenBank Accession Number:

NM\_007932.2

GeneID (NCBI):

100ug, Concentration: 1 mg/ml by 13805

**UNIPROT ID:** 

Nanodrop: Source: Q63961 Rabbit Full Name: Isotype: endoglin

Calculated MW:

70kDa

Observed MW: 85-90 kDa

**Applications** 

**Tested Applications:** 

WB, IHC, IF/ICC, Indirect ELISA

Species Specificity:

mouse

IgG

**Background Information** 

Endoglin (ENG, CD105) is a homodimeric cell membrane glycoprotein of 180 kDa, composed of disulphide-linked subunits of 90-95 kDa. Endoglin is a proliferation-associated and hypoxia-inducible protein mainly expressed on vascular endothelial cells. It acts as an accessory receptor for transforming growth factor beta  $(TFG-\beta)$  and is involved in vascular development and remodeling. The important role of Endoglin in angiogenesis and in tumor progression makes it an ideal target for antiangiogenic therapy and a good marker for tumor prognosis. (PMID: 1326540; 14528280; 21737653; 12773481)


Storage

Store at -80°C. Storage Buffer: PBS Only


in USA), or 1(312) 455-8498 (outside USA)

W: ptglab.com


## Selected Validation Data




bEnd.3 cells were subjected to SDS PAGE followed by western blot with 84807-4-RR (Endoglin/CD105 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 84807-4-PBS in a different storage buffer formulation.




Immunofluorescent analysis of (4% PFA) fixed bEnd.3 cells using Endoglin/CD105 antibody (84807-4-RR, Clone: 242305G3) at dilution of 1:400 and Coralite@488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-Phalloidin (red). This data was developed using the same antibody clone with 84807-4-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded mouse kidney tissue slide using 84807-4-RR (Endoglin/CD105 antibody) at dilution of 1:800 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 84807-4-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded mouse kidney tissue slide using 84807-4-RR (Endoglin/CD105 antibody) at dilution of 1:800 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 84807-4-PBS in a different storage buffer formulation.



Biolayer interferometry (BLL) kinetic assays of 84807-4-RR against Mouse Endoglin/CD105 were performed. The affinity constant is 0.888 nM.