## For Research Use Only

## P62/SQSTM1 Recombinant antibody

Catalog Number:84826-1-RR



**Basic Information** 

Catalog Number: GenBank Accession Number:

84826-1-RR NM 011018 GeneID (NCBI):

100ul , Concentration: 1000  $\mu g/ml$  by 18412 Nanodrop: **UNIPROT ID:** 

Q64337 Rabbit Full Name:

Isotype: sequestosome 1 IgG Calculated MW:

Immunogen Catalog Number: 48 kDa

AG35064 Observed MW:

62 kDa

**Purification Method:** 

Protein A purification

CloneNo.: 241992C4

Recommended Dilutions:

WB 1:5000-1:50000

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:200-1:800 IF/ICC 1:250-1:1000

**Applications** 

**Tested Applications:** 

WB, IHC, IF/ICC, FC (Intra), IP, ELISA

Species Specificity:

human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

WB: NIH/3T3 cells, HeLa cells, mouse brain tissue, HepG2 cells, Jurkat cells, Raji cells, HEK-293T cells, MCF-7 cells, mouse liver tissue, rat brain tissue

IP: NIH/3T3 cells.

IHC: mouse brain tissue, IF/ICC: NIH/3T3 cells,

**Background Information** 

Sequestosome 1 (SQSTM1/p62) is a multifunctional adaptor protein implicated in selective autophagy, cell signaling pathways, and tumorigenesis. p62 has been implicated in shuttling ubiquitinated and aggregated proteins for autophagic degradation. p62 is degraded during the autophagic process, which makes its intracellular level a marker for autophagy progression. p62 is at the cross-roads of several signaling pathways including Ras/ Raf/ MAPK and NFkB and plays important role in cancer.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol, pH7.3


Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


## Selected Validation Data




NIH/3T3 cells were subjected to SDS PAGE followed by western blot with 84826-1-RR (P62/SQSTM1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 84826-1-RR (P62/SQSTM1 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.




Various lysates were subjected to SDS PAGE followed by western blot with 84826-1-RR (P62/SQSTM1 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.




IP result of anti-P62/SQSTM1 (IP:84826-1-RR, 4ug; Detection:84826-1-RR 1:4000) with NIH/3T3 cells lysate 1920 ug.




Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 84826-1-RR (P62/SQSTM1 antibody) at dilution of 1:400 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 84826-1-RR (P62/SQSTM1 antibody) at dilution of 1:400 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed NIH/3T3 cells using P62/SQSTM1 antibody (84826-1-RR, Clone: 241992C4) at dilution of 1:500 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-Phalloidin (red).



1x10^6 NIH/3T3 cells were intracellularly stained with 0.25 ug P62/SQSTM1 Recombinant antibody (84826-1-RR, Clone:241992C4) and Coralite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2) (red), or 0.25 ug Rabbit IgG Isotype Control RecAb (98136-1-RR, Clone: 240953C9) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



Biolayer interferometry (BLL) kinetic assays of 84826-1-RR against Mouse P62,SQSTM1 were performed. The affinity constant is 0.203 nM.