## For Research Use Only

## CD3 Gamma Recombinant antibody

Catalog Number:84836-1-RR



**Purification Method:** 

**Basic Information** 

Catalog Number: GenBank Accession Number:

84836-1-RR BC113830 Protein A purfication

GeneID (NCBI): CloneNo.: 100ul , Concentration: 1000  $\mu g/ml$  by 917 241849E4

Nanodrop: **UNIPROT ID:** Recommended Dilutions: Source: P09693 WB 1:5000-1:50000 Rabbit IHC 1:500-1:2000 Full Name: IF/ICC 1:200-1:800

Isotype: CD3g molecule, gamma (CD3-TCR IgG

complex)

Calculated MW: 182 aa, 20 kDa Observed MW: 20-25 kDa

**Applications** 

**Tested Applications:** WB, IHC, IF/ICC, ELISA

Species Specificity:

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: Jurkat cells, MOLT-4 cells IHC: human tonsillitis tissue,

IF/ICC: Jurkat cells,

## **Background Information**

CD3 is a complex of proteins that directly associates with the T cell receptor (TCR). The TCR/CD3 complex of Tlymphocytes consists of either a TCR alpha/beta or TCR gamma/delta heterodimer coexpressed at the cell surface with the invariant subunits of CD3 labeled gamma, delta, epsilon, zeta, and eta. The TCR recognizes antigens bound to major histocompatibility complex (MHC) molecules. TCR-mediated peptide-MHC recognition is transmitted to the CD3 complex, leading to the intracellular signal transduction. CD3 is considered to be a pan-T cell marker for detection of normal and neoplastic T cells.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


\*\*\* 20ul sizes contain 0.1% BSA

in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## **Selected Validation Data**




Various lysates were subjected to SDS PAGE followed by western blot with 84836-1-RR (CD3 Gamma antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 84836-1-RR (CD3 Gamma antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 84836-1-RR (CD3 Gamma antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed Jurkat cells using CD3 Gamma antibody (84836-1-RR, Clone: 241849E4) at dilution of 1:400 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2).