For Research Use Only

## HCN1 Recombinant antibody, PBS Only

Catalog Number:84845-3-PBS



**Basic Information** 

Catalog Number:

GenBank Accession Number:

**Purification Method:** 

84845-3-PBS

NM\_021072 GeneID (NCBI): Protein A purfication

CloneNo.:

100ug, Concentration: 1 mg/ml by Nanodrop;

348980 **UNIPROT ID:**  242254A1

Source: Rabbit

060741

Isotype:

IgG

Full Name:

 $hyperpolarization\,activated\,cyclic$ nucleotide-gated potassium channel

Calculated MW:

99 kDa

Observed MW:

99-120 kDa

**Applications** 

**Tested Applications:** 

WB, IHC, Indirect ELISA

Species Specificity:

human, mouse

## **Background Information**

 $HCN1, also \ named \ as \ BCNG1, belongs \ to \ the \ potassium \ channel \ HCN \ family. \ It \ hyperpolarization-activated \ ion$ channel exhibiting weak selectivity for potassium over sodium ions. HCN1 contributes to the native pacemaker currents in heart (If) and in neurons (Ih). It is activated by cAMP, and at 10-100 times higher concentrations, also by cGMP. HCN1 may mediate responses to sour stimuli.

Storage


Storage:

Store at -80°C.

Storage Buffer:

PBS Only

## **Selected Validation Data**



Boiled and unboiled mouse brain tissue lysates were subjected to SDS PAGE followed by western blot with 84845-3-RR (HCN1 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 84845-3-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded mouse cerebellum tissue slide using 84845-3-RR (HCN1 antibody) at dilution of 1:400 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 84845-3-PBS in a different storage buffer formulation.