For Research Use Only

Phospho-EGFR (Tyr1197) Recombinant antibody

Catalog Number:84906-1-RR





Basic Information	Catalog Number: 84906-1-RR	GenBank Accession Number: BC094761	Purification Method: Protein A purfication
	Size: 100ul , Concentration: 1000 µg/ml by Nanodrop; Source: Rabbit Isotype: IgG	GeneID (NCBI): 1956	CloneNo.: 241826G10
		UNIPROT ID: P00533	Recommended Dilutions: WB 1:1000-1:4000
		Full Name: epidermal growth factor receptor (erythroblastic leukemia viral (v-erb- b) oncogene homolog, avian) Calculated MW: 1210 aa, 134 kDa	
		Applications	Tested Applications:
WB, FC (Intra), ELISA Species Specificity: human			
Background Information	Epidermal Growth Factor Receptor (EGFR) is one of the receptor tyrosine kinases (RTKs), members of the ErbB/HER family which consists of ErbB1 (EGFR or HER1), ErbB2 (HER2 or Neu), ErbB3 (HER3), and ErbB4 (HER4). The EGFR autophosphorylation sites Tyr-992 and Tyr-1173 (also known as Tyr 1197) have been shown to play a critical role in the activation of the MAPK cascade following EGF stimulation. (PMID: 15708852)		
Storage	Storage: Store at -20°C. Stable for one year after shipment. Storage Buffer: PBS with 0.02% sodium azide and 50% glycerol pH 7.3.		
*** 20ul sizes contain 0.1% BSA	Aliquoting is unnecessary for -20°C s		

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)E: proteintech@ptglab.comW: ptglab.comW: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





Non-treated A431 cells and EGF treated A431 cells were subjected to SDS PAGE followed by western blot with 84906-1-RR (Phospho-EGFR (Tyr1197) antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours. The membrane was stripped and re-blotted with Alpha Tubulin (66031-1-lg) antibody as a loading control. 1x10^6 A431 cells untreated (dashed lines) or treated with EGF which intracellularly stained with 0.25 ug Phospho-EGFR (Tyr1197) Recombinant Antibody (84906-1-RR, Clone:241826G10) and PE-Conjugated Goat Anti-Rabbit IgG(H+L)(red), or 0.25 ug Rabbit IgG Isotype Control Recombinant Antibody (98136-1-RR, Clone: 240953C9) (blue). Cells were fixed with 4% PFA and permeabilized with 90% MeOH.