

For Research Use Only

Tri-Methyl-Histone H3 (Lys4) Recombinant monoclonal antibody

Catalog Number: 84908-3-RR

5 Publications



Basic Information

Catalog Number:

84908-3-RR

Size:

100ul, Concentration: 1000 µg/ml by Nanodrop;

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

BC066245

GeneID (NCBI):

8350

UNIPROT ID:

P68431

Full Name:

histone cluster 1, H3a

Observed MW:

15 kDa

Purification Method:

Protein A purification

CloneNo.:

241942H7

Recommended Dilutions:

WB: 1:5000-1:50000

Dot Blot: 1:500-1:2000

ChIP-qPCR: 1:10-1:100

Applications

Tested Applications:

WB, Dot Blot, ELISA, ChIP-qPCR

Cited Applications:

WB, ChIP

Species Specificity:

human, mouse, rat

Cited Species:

human, mouse, rat

Positive Controls:

WB : HeLa cells, K-562 cells, HEK-293T cells, MCF-7 cells, NIH/3T3 cells, HSC-T6 cells

Dot Blot : / /,

ChIP-qPCR : HeLa cells,

Background Information

Histones are small, highly basic proteins that consist of a globular domain with unstructured N- and C-terminal tails protruding from the main structure. Histone H3 is one of the five main histones that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. In addition to their role in DNA compartmentalization, histones also play crucial roles in various biologic processes, including gene expression and regulation, DNA repair, chromatin condensation, cell cycle progression, chromosome segregation, and apoptosis. The ability of histones to regulate chromatin dynamics primarily originates from various posttranslational modifications carried out by histone-modifying enzymes.

Notable Publications

Author	Pubmed ID	Journal	Application
Peng Cui	42449933	Int J Mol Sci	ChIP
Hangjie Ying	42229717	J Adv Res	WB
Yan Zhang	41796727	Biochem Pharmacol	ChIP, WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Dot blot analysis was used to confirm the specificity of 84908-3-RR Tri-Methyl-Histone H3 (Lys4) antibody. peptides were spotted onto NC and probed with antibody at 1 µg/mL. The amount of peptide (µg/mL) spotted is indicated next to each row.



Chromatin was prepared from HeLa cells. Cells were fixed with formaldehyde for 10 minutes. The ChIP was performed with 15 µg of cross-linked chromatin, 5 µg of Tri-Methyl-Histone H3 (Lys4) (84908-3-RR) or 5 µg of Normal Rabbit IgG (98136-1-RR), and 20 µl of Protein A Magarose Beads. The immunoprecipitated DNA was quantified by real-time PCR.



Various lysates were subjected to SDS PAGE followed by western blot with 84908-3-RR (Tri-Methyl-Histone H3 (Lys4) antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 84908-3-RR (Tri-Methyl-Histone H3 (Lys4) antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.