For Research Use Only

## TYRO3 Recombinant antibody

Catalog Number:84929-6-RR




Basic Information	Catalog Number: 84929-6-RR	GenBank Accession Number: BC049368	Purification Method: Protein A purfication			
	Size: 100ul , Concentration: 1000 µg/ml by Nanodrop; Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG29527	UNIPROT ID: Q06418 Full Name:	CloneNo.: 242477G1 Recommended Dilutions: WB 1:5000-1:50000 IHC 1:125-1:500			
				TYRO 3 protein tyrosine kinase Calculated MW: 97 kDa Observed MW: 130 kDa		
						Applications
		Species Specificity: human, mouse, rat	WB : mouse brain tissue, IHC : mouse brain tissue,			
		Note-IHC: suggested antigen r TE buffer pH 9.0; (*) Alternativ retrieval may be performed w buffer pH 6.0	vely, antigen			
Background Information	TYRO3 has a structure includes a cytoplasmic domain and an extracellular ligand binding region. And ligand binding at the cell surface induces dimerization and autophosphorylation of TYRO3 on its intracellular domain that provides docking sites for downstream signaling molecules. TYRO3 is associated with many physiological processes including cell survival, migration and differentiation. It has been reported that TYRO3 can be expressed in various regions of the mammalian central nervous system. The molecular mass of TYRO3 is 130 kDa, the apparent molecular mass is higher due to glycosylation.((PMID: 8873131, 23354874, 21539887)					
Storage	Storage: Store at -20°C. Stable for one year aft Storage Buffer: PBS with 0.02% sodium azide and 50					
*** 20ul sizes contain 0.1% BSA	Aliquoting is unnecessary for -20 $^{\circ}$ C s	torage				

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data





Various lysates were subjected to SDS PAGE followed by western blot with 84929-6-RR (TYRO3 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.

Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 84929-6-RR (TYRO3 antibody) at dilution of 1:250 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 84929-6-RR (TYRO3 antibody) at dilution of 1:250 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Biolayer interferometry (BLl) kinetic assays of 84929-6-RR against Human TYRO3 were performed. The affinity constant is 0.133 nM.