For Research Use Only

## Caspase 9 Recombinant antibody

Catalog Number:84949-4-RR




Basic Information	Catalog Number: 84949-4-RR	GenBank Accession Number: BC002452	Purification Method: Protein A purification
	Size: 100ul , Concentration: 1000 µg/ml by Nanodrop; Source: Rabbit Isotype: IgG	GeneID (NCBI): 842	CloneNo.: 242452F12
		UNIPROT ID: P55211	Recommended Dilutions: WB 1:1000-1:4000
		Full Name:	
		caspase 9, apoptosis-related cysteine peptidase Calculated MW: 46 kDa	
		Applications	Tested Applications: WB, ELISA Species Specificity: human
WB : Jurka	at cells,		
Background Information	Caspase 9, apoptosis-related cysteine protease (CASP9, synonyms: MCH6, APAF3, APAF-3, ICE-LAP6, CASPASE-9c) is a member of the cysteine-aspartic acid protease (caspase) family. Sequential activation of caspases plays a central role in the execution-phase of cell apoptosis. Caspases exist as inactive proenzymes which undergo proteolytic processing at conserved aspartic residues to produce 2 subunits, large and small, that dimerize to form the active enzyme. Caspase 9 is processed by APAF1; this step is thought to be one of the earliest in the caspase activation cascade. 10380-1-AP can recognize the pre- and cleaved- caspase 9. In recent years, the localization of caspase 9 was a focus of interest. Beside its cytoplasmic distribution, a very extensive localization study was done on rat brain tissue, where caspase 9 was found located predominantly in the nucleus and to a lesser extend in the cytoplasm [PMID: 15541731].		
	cascade. 10380-1-AP can recognize th was a focus of interest. Beside its cyto tissue, where caspase 9 was found loo	e pre- and cleaved- caspase 9. In poplasmic distribution, a very exter	e of the earliest in the caspase activation recent years, the localization of caspase 9 nsive localization study was done on rat brain
Storage	cascade. 10380-1-AP can recognize th was a focus of interest. Beside its cyto tissue, where caspase 9 was found loo	e pre- and cleaved- caspase 9. In p pplasmic distribution, a very exter ated predominantly in the nucleu er shipment. % glycerol, pH7.3	e of the earliest in the caspase activation recent years, the localization of caspase 9 1sive localization study was done on rat brain

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free<br/>in USA), or 1(312) 455-8498 (outside USA)E: proteintech@ptglab.comW: ptglab.comW: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 84949-4-RR (Caspase 9 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.