For Research Use Only

PBK/SPK Recombinant antibody, PBS Only (Capture)

Catalog Number:84970-1-PBS



Purification Method:

Protein A purification

CloneNo.:

242545A11

Basic Information

Catalog Number: GenBank Accession Number:

84970-1-PBS BC015191
Size: GeneID (NCBI):

100ug, Concentration: 1 mg/ml by 55872

Nanodrop; UNIPROT ID:
Source: Q96KB5
Rabbit Full Name:

Isotype: PDZ binding kinase
IgG Calculated MW:
Immunogen Catalog Number: 322 aa, 36 kDa
AG9033 Observed MW:

36-40 kDa

Applications

Tested Applications:

WB, IF/ICC, FC (Intra), Cytometric bead array, Sandwich ELISA, Indirect ELISA, Sample test

Species Specificity: human, mouse, rat

Product Information

84970-1-PBS targets PBK/SPK as part of a matched antibody pair:

MP01693-1: 84970-1-PBS capture and 84970-4-PBS detection (validated in Cytometric bead array, Sandwich ELISA)

Unconjugated rabbit recombinant monoclonal antibody in PBS only (BSA and azide free) storage buffer at a concentration of 1 mg/mL, ready for conjugation. Created using Proteintech's proprietary in-house recombinant technology. Recombinant production enables unrivalled batch-to-batch consistency, easy scale-up, and future security of supply.

This conjugation ready format makes antibodies ideal for use in many applications including: ELISAs, multiplex assays requiring matched pairs, mass cytometry, and multiplex imaging applications. Antibody use should be optimized by the end user for each application and assay.


Background Information

PBK(PDZ-binding kinase) is also named as TOPK, CT84, Nori-3, SPK and belongs to the MAP kinase kinase subfamily. PBK may have a role in the regulation of cellular proliferation and progression of the cell cycle. It has a characteristic cdc2ycyclin B phosphorylation site(SyT-P-X-KyR) at its N terminus, which is conserved across species, and it is phosphorylated in a cell cycle-dependent manner at mitosis, and that this phosphorylation is required for its activation(PMID:10779557).


Storage

Storage: Store at -80°C. Storage Buffer: PBS only, pH7.3


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 84970-1-RR (PBK antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 84970-1-PBS in a different storage buffer formulation.




1x10^6 HepG2 cells were intracellularly stained with 0.25 ug PBK/SPK Recombinant antibody (84970-1-RR, Clone:242545A11) and Coralite@488-Conjugated Goat Anti-Rabbit 1gG(H+L) (SA00013-2)(red), or 0.25 ug Rabbit 1gG Isotype Control RecAb (98136-1-RR, Clone: 240953C9) (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011). This data was developed using the same antibody clone with 84970-1-PBS in a




Sandwich ELISA standard curve of MP01693-1, Human PBK Recombinant Matched Antibody Pair -PBS only. 84970-1-PBS was coated to a plate as the capture antibody and incubated with serial dilutions of standard Ag9033. 84970-4-PBS was HRP conjugated as the detection antibody. Range: 0.781-50 ng/mL



The mean PBK concentration was determined to be 342.86 ng/mL in Hela cell extract based on a 1.30 mg/mL extract load and 1,243.51 ng/mL in Jurkat cell extract based on a 2.00 mg/mL extract load.



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using PBK/SPK antibody (84970-1-RR, Clone: 242545A11) at dilution of 1:1000 and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-Phalloidin (red). This data was developed using the same antibody clone with 84970-1-PBS in a different storage buffer formulation.



Cytometric bead array standard curve of MP01693-1, PBK Recombinant Matched Antibody Pair, PBS Only. Capture antibody: 84970-1-PBS. Detection antibody: 84970-4-PBS. Standard: Ag9033. Range: 0.781-100 ng/mL