

For Research Use Only

ADAR1 Recombinant antibody

Catalog Number: 85009-1-RR



Basic Information

Catalog Number: 85009-1-RR	GenBank Accession Number: BC038227	Purification Method: Protein A purification
Size: 100ul , Concentration: 1000 µg/ml by Nanodrop;	GeneID (NCBI): 103	CloneNo.: 241595E8
Source: Rabbit	UNIPROT ID: P55265	Recommended Dilutions: WB 1:500-1:2000
Isotype: IgG	Full Name: adenosine deaminase, RNA-specific	
Immunogen Catalog Number: AG5609	Calculated MW: 136 kDa	
	Observed MW: 110 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB : HepG2 cells, HeLa cells, HEK-293T cells, A431 cells
Species Specificity: human	

Background Information

ADAR1 is also named as ADAR1, DSRAD, G1P1, IFI4. It convert selected adenosine residues into inosine in substrate RNAs containing a relatively short dsRNA region(PMID:15556947). The human ADAR1 gene specifies two size forms of RNA-specific adenosine deaminase, an IFN inducible 150 kDa protein and a constitutively expressed N-terminally truncated 110 kDa protein, encoded by transcripts with alternative exon 1 structures that initiate from different promoters(PMID:11111054). It has 5 isoforms produced by alternative promoter usage and alternative splicing. Defects in ADAR are a cause of dyschromatosis symmetrical hereditaria (DSH).ADAR1 can form respective homodimers, and this association is essential for its enzymatic activities(PMID:17428802).

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 85009-1-RR (ADAR1 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.