

For Research Use Only

BAG2 Recombinant antibody, PBS Only

Catalog Number: 85063-4-PBS



Basic Information

Catalog Number: 85063-4-PBS	GenBank Accession Number: NM_004282	Purification Method: Protein A purification
Size: 100ug, Concentration: 1 mg/ml by Nanodrop;	GeneID (NCBI): 9532	CloneNo.: 242717G4
Source: Rabbit	UNIPROT ID: O95816	
Isotype: IgG	Full Name: BCL2-associated athanogene 2	
Immunogen Catalog Number: AG30873	Calculated MW: 24 kDa	
	Observed MW: 25 kDa	

Applications

Tested Applications:
WB, Indirect ELISA

Species Specificity:
human, mouse, rat

Background Information

BAG2 (BAG family molecular chaperone regulator 2) is one of six proteins in mammals that contain the BAG domain, which belongs to the BAG (Bcl-2-associated athanogene) family. BAG2 has been described as a negative regulator of the chaperone-associated ubiquitin ligase C terminus of Hsc70-interacting protein (CHIP) that participates in the ubiquitin-mediated proteasomal degradation of misfolded substrate proteins (PMID: 16207813). BAG2 is widely expressed in human tissues, including brown adipose, heart and lung tissue, as well as in various types of tumor cells, including renal cell carcinoma, glioblastoma and thyroid carcinoma cells (PMID: 28536620). BAG2 overexpression is associated with poor prognosis in patients and mtp53 accumulation in tumors (PMID: 26271008).

Storage


Storage:
Store at -80°C.

Storage Buffer:
PBS only, pH7.3


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 85063-4-RR (BAG2 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 85063-4-PBS in a different storage buffer formulation.



Various lysates were subjected to SDS PAGE followed by western blot with 85063-4-RR (BAG2 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 85063-4-PBS in a different storage buffer formulation.



Biolayer interferometry (BLI) kinetic assays of 85063-4-RR against Human BAG2 were performed. The affinity constant is 0.644 nM.