For Research Use Only

Arrestin C Recombinant antibody

Catalog Number:85067-5-RR




Basic Information	Catalog Number: 85067-5-RR	GenBank Accession Number: BC012096	Purification Method: Protein A purification
	Size: 100ul , Concentration: 1000 µg/ml by Nanodrop; Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG1580	GenelD (NCBI): 407	CloneNo.: 242751B12
		UNIPROT ID: P36575 Full Name: arrestin 3, retinal (X-arrestin) Calculated MW: 43 kDa	Recommended Dilutions: WB 1:2000-1:10000 IF-P 1:500-1:2000
		Observed MW: 47-50 kDa	
Applications	Tested Applications: WB, IF-P, ELISA Species Specificity: human, mouse, rat	Positive Controls: WB : mouse eye tissue, rat eye tissues IF-P : mouse eye tissue,	
Background Information	Arrestin C, also known as Arrestin 3 or Retinal Cone Arrestin, is a protein encoded by the ARR3 gene. It belongs to the arrestin family, which plays a crucial role in regulating G-protein-coupled receptor (GPCR) signaling and trafficking. Arrestin C is composed of two major domains: the N-domain and the C-domain, connected by a hinge region. These domains form a structure resembling two clamshells placed end-to-end. The C-terminal tail (C-tail) of Arrestin C interacts extensively with the N-domain, stabilizing its basal conformation. Arrestin C is predominantly expressed in cone photoreceptors and pinealocytes in the retina. It is involved in the shut-off mechanisms associated with high-acuity color vision by binding to phosphorylated and activated opsins, thereby inhibiting their ability to interact with transducin.		
Storage	Storage: Store at -20°C. Stable for one year after Storage Buffer: PBS with 0.02% sodium azide and 50° Aliquoting is unnecessary for -20°C st	% glycerol pH 7.3.	
	Auquoting is unnecessary 101-20 C S	loidge	


For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)E: proteintech@ptglab.comW: ptglab.comW: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 85067-5-RR (ARR3 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded mouse eye tissue using Arrestin C antibody (85067-5-RR, Clone: 242751B12) at dilution of 1:1000 and CoraLite@488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded mouse eye tissue using Arrestin C antibody (85067-5-RR, Clone: 242751B12) at dilution of 1:1000 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Biolayer interferometry (BLL) kinetic assays of 85067-5-RR against Human Arrestin C were performed. The affinity constant is below 1 pM.