

For Research Use Only

VLDLR Recombinant antibody

Catalog Number: 85097-1-RR



Basic Information

Catalog Number: 85097-1-RR	GenBank Accession Number: NM_003383	Purification Method: Protein A purification
Size: 100ul , Concentration: 1000 µg/ml by Nanodrop;	GeneID (NCBI): 7436	CloneNo.: 242385H3
Source: Rabbit	UNIPROT ID: P98155	Recommended Dilutions: WB: 1:5000-1:50000
Isotype: IgG	Full Name: very low density lipoprotein receptor	FC (Intra): 0.25 ug per 10 ⁶ cells in a 100 µl suspension
	Calculated MW: 96 kDa	
	Observed MW: 120 kDa	

Applications

Tested Applications: WB, FC (Intra), ELISA	Positive Controls:
Species Specificity: human	WB : MCF-7 cells, THP-1 cells
	FC (Intra) : THP-1 cells,

Background Information

The very low-density lipoprotein receptor (VLDLR) is a member of the LDL receptor family with a complex molecular weight profile due to post-translational modifications. The calculated molecular weight of the core protein is approximately 84-86 kDa, however, VLDLR undergoes extensive glycosylation during processing, resulting in higher apparent molecular weights observed in SDS-PAGE analyses. Multiple studies report that glycosylated VLDLR migrates as broad bands between 90-180 kDa under reducing conditions, with the most commonly observed forms at approximately 96 kDa (core glycosylated form), 115-135 kDa (mature fully glycosylated form), and 150-180 kDa (hyperglycosylated or dimeric forms).

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol, pH7.3
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 85097-1-RR (VLDLR antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



1x10⁶ THP-1 cells were intracellularly stained with 0.25 ug VLDLR Recombinant Antibody (85097-1-RR, Clone:242385H3) and APC-Conjugated Goat Anti-Rabbit IgG(H+L)(red), or 0.25 ug Rabbit IgG Isotype Control Recombinant Antibody (98136-1-RR, Clone: 240953C9) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).