For Research Use Only

TORC2 Recombinant antibody

Catalog Number:85150-4-RR




Basic Information	Catalog Number: 85150-4-RR	GenBank Accession Number: BC053562	Purification Method: Protein A purification
	Size: 100ul , Concentration: 1000 µg/ml by Nanodrop; Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG3167	Genel D (NCBI): 200186	CloneNo.: 242808B6
		UNIPROT ID: Q53ETO Full Name: CREB regulated transcription coactivator 2	Recommended Dilutions: WB 1:5000-1:50000
		Observed MW: 80 kDa	
		Applications	Tested Applications:
WB, FC (Intra), ELISA Species Specificity: human	WB : HEK-293 cells, Jurkat cells, Raji cells, A-673 cells		
Background Information	CRTC2, also named as TORC2, belongs to the TORC family. It is a transcriptional coactivator for CREB1 which activates transcription through both consensus and variant cAMP response element (CRE) sites. It acts as a coactivator, in the SIK/TORC signaling pathway, being active when dephosphorylated and acts independently of CREB1 'Ser-133' phosphorylation. CRTC2 enhances the interaction of CREB1 with TAF4. Ir regulates gluconeogenesis as a component of the LKB1/AMPK/TORC2 signaling pathway. CRTC2 regulates the expression of specific genes such as the steroidogenic gene, StAR. TORC2 was recently shown to be an important regulator of gluconeogenesis in the livers of mammals. It is one of the other key regulators of CRE-dependent MIE gene expression in NT2 cells. This regulation is linked to VIP-induced TORC2 dephosphorylation and translocation to the nucleus. (PMID: 19369332, 20504934).		
Storage	Storage: Store at -20°C. Stable for one year aft Storage Buffer:	•	
*** 20ul sizes contain 0.1% BSA	PBS with 0.02% sodium azide and 50 Aliquoting is unnecessary for -20° C s	•••	


For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)E: proteintech@ptglab.comW: ptglab.comW: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 85150-4-RR (CRTC2,TORC2 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



1x10^6 HeLa cells were intracellularly stained with 0.25 ug TORC2 Recombinant antibody (85150-4-RR, Clone:242808B6) and CoraLite®488-Conjugated Goat Anti-Rabbit 1gG(H+L) (5A00013-2)(red), or 0.25 ug Rabbit 1gG Isotype Control RecAb (98136-1-RR, Clone: 240953C9) (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).



Biolayer interferometry (BLL) kinetic assays of 85150-4-RR against Human TORC2 were performed. The affinity constant is 0.422 nM.