

For Research Use Only

Cleaved PARP1 (Asp214), p25 Recombinant monoclonal antibody, PBS Only

Catalog Number: 85175-6-PBS



Basic Information

Catalog Number: 85175-6-PBS	GenBank Accession Number: BC037545	Purification Method: Protein A purification
Size: 100ug , Concentration: 1 mg/ml by Nanodrop;	GeneID (NCBI): 142	CloneNo.: 253204E4
Source: Rabbit	UNIPROT ID: P09874	
Isotype: IgG	Full Name: poly (ADP-ribose) polymerase 1	
	Calculated MW: 1014 aa, 113 kDa	
	Observed MW: 25 kDa	

Applications

Tested Applications:
WB, Indirect ELISA

Species Specificity:
human

Background Information

PARP1 (poly(ADP-ribose) polymerase 1) is a nuclear enzyme catalyzing the poly(ADP-ribosyl)ation of many key proteins in vivo. The normal function of PARP1 is the routine repair of DNA damage. Activated by DNA strand breaks, the PARP1 is cleaved into an 85 to 89-kDa COOH-terminal fragment and a 24 kDa NH2-terminal peptide by caspases during the apoptotic process. The appearance of PARP fragments is commonly considered an important biomarker of apoptosis. In addition to caspases, other proteases like calpains, cathepsins, granzymes, and matrix metalloproteinases (MMPs) have also been reported to cleave PARP1 and give rise to fragments ranging from 42-89 kDa. This antibody only recognizes the smaller cleaved form of PARP1 (around 25 kDa) but not full-length PARP1.

Storage


Storage:
Store at -80°C.

Storage Buffer:
PBS only, pH7.3


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Non-treated Jurkat cells and Staurosporine-treated Jurkat cells were subjected to SDS PAGE followed by western blot with 85175-6-RR (Cleaved PARP1 (Asp214), p25 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and re-blotted with Beta-Actin (66009-1-Ig) antibody as a loading control. This data was developed using the same antibody clone with 85175-6-PBS in a different storage buffer formulation.



Non-treated HeLa cells and Staurosporine-treated HeLa cells were subjected to SDS PAGE followed by western blot with 85175-6-RR (Cleaved PARP1 (Asp214), p25 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and re-blotted with Beta-Actin (66009-1-Ig) antibody as a loading control. This data was developed using the same antibody clone with 85175-6-PBS in a different storage buffer formulation.