For Research Use Only

TRPC4 Recombinant antibody

Catalog Number:85334-2-RR



Purification Method:

Protein A purification

Recommended Dilutions:

WB 1:2000-1:10000

CloneNo.:

242444F8

Basic Information

Catalog Number: GenBank Accession Number:

85334-2-RR BC104725 GeneID (NCBI): Size:

100ul , Concentration: 1000 $\mu g/ml$ by 7223 Nanodrop: **UNIPROT ID:** Q9UBN4

Rabbit Full Name:

Isotype: transient receptor potential cation channel, subfamily C, member 4 IgG

Immunogen Catalog Number: Calculated MW: 977 aa, 112 kDa AG15698

> Observed MW: 140 kDa

Applications

Tested Applications:

WB, ELISA

Species Specificity:

human

Positive Controls:

WB: A549 cells, HEK-293 cells, SH-SY5Y cells

Background Information

TRPC4 (Transient Receptor Potential Paradigm 4) is a non-selective calcium channel protein widely found in the nervous system and cardiovascular system. TRPC4 regulates intracellular calcium levels through the activation of $signaling\ pathways\ mediated\ by\ Gq/11\ and\ Gi/o-coupled\ receptors, and\ is\ involved\ in\ neurotransmission,\ neuronal$ excitability, and vascular endothelial cell function. In the nervous system, TRPC4 is associated with neuropsychiatric disorders such as depression and anxiety, and its inhibitors show antidepressant and anxiolytic potential. In the cardiovascular system, TRPC4 affects vascular permeability and diastolic function by regulating $calcium\ ion\ inward\ flow\ and\ plays\ an\ important\ role\ in\ hypoxia-induced\ pulmonary\ hypertension.\ In\ addition,$ TRPC4 is involved in thermore gulation, sensing changes in internal temperature to regulate the drop in body and the property of the properttemperature. aberrant expression of TRPC4 is associated with a variety of diseases, making it a potential therapeutic target!

Storage

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 85334-2-RR (TRPC4 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.