For Research Use Only

Cleaved Caspase 6/P18 Recombinant antibody

Catalog Number:85337-1-RR




Basic Information	Catalog Number: 85337-1-RR	GenBank Accession Number: BC 000305	Purification Method: Protein A purification
	Size: 100ul , Concentration: 1000 µg/ml by Nanodrop; Source: Rabbit Isotype: IgG	GenelD (NCBI): 839	CloneNo.: 242384E6
		UNIPROT ID: P55212 Full Name:	Recommended Dilutions: WB 1:1000-1:4000
		caspase 6, apoptosis-related cysteine peptidase Calculated MW: 33 kDa, 22 kDa	
		Applications	Tested Applications: WB, ELISA Species Specificity: human
WB : Stau	rosporine treated Jurkat cells,		
Background Information	Caspase-6 belongs to caspase family of cysteinyl-aspartate specific proteases.Precursor of CASP6 produces two subunits, large (18kDa) and small (16kDa) that dimerize. It cleaves poly(ADP-ribose) polymerase, as well as lamins and is involved in the activation cascade of caspases responsible for apoptosis execution. Researches showed that CASP6 could be an early instigator of neuronal dysfunction and regulates B cell activation and differentiation into plasma cells by modifying cell cycle entry. IRAK3 is an important target for CASP6. It can reveal five bands of 28, 32, 36, 49, and 64 kDa in human neurons and fetal brain in western blot, the 32 and 28 kDa bands represent procaspase-6 and pro-arm caspase-6. Procaspase-6 is more abundant than pro-arm caspase-6 in adult tissue, whereas pro-arm caspase-6 is more abundant than pro-caspase-6 in fetal brain and cultured neurons. This antibody is specific to Caspase 6 P18.		
Storage	Storage: Store at -20°C. Stable for one year after shipment. Storage Buffer: PBS with 0.02% sodium azide and 50% glycerol, pH7.3		
*** 20ul sizes contain 0.1% BSA	Aliquoting is unnecessary for -20 $^{\circ}$ C s	torage	

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)E: proteintech@ptglab.comW: ptglab.comW: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Staurosporine treated Jurkat cells were subjected to SDS PAGE followed by western blot with 85337-1-RR (Cleaved Caspase 6/P18 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.