For Research Use Only

TRIM33 Recombinant antibody, PBS Only

Catalog Number:85364-3-PBS



Basic Information

Catalog Number:

85364-3-PBS

Size:

Source:

NM_015906 GeneID (NCBI):

GenBank Accession Number:

100ug, Concentration: 1 mg/ml by Nanodrop:

51592 **UNIPROT ID:** Q9UPN9 Full Name:

Rabbit Isotype: IgG

tripartite motif-containing 33 Calculated MW:

122 kDa Observed MW: 140-150 kDa

Purification Method:

Protein A purification CloneNo.:

242728H2

Applications

Tested Applications: WB, IF/ICC, Indirect ELISA Species Specificity:

human


Background Information

TRIM33, also named KIAA1113, RFG7, and TIF1G, belongs to the TRIM/RBCC family. It acts as an E3 ubiquitin-protein ligase. TRIM33 promotes SMAD4 ubiquitination, nuclear exclusion, and degradation via the ubiquitin-proteasome pathway. It may act as a transcriptional repressor. It inhibits the transcriptional response to TGF-beta/BMP signaling cascade. TRIM33 plays a role in the control of cell proliferation. Its association with SMAD2 and SMAD3, TRIM33 $stimulates\ erythroid\ differentiation\ of\ hematopoietic\ stem/progenitor.\ Defects\ in\ TRIM33\ are\ a\ cause\ of\ thyroid\ differentiation\ of\ hematopoietic\ stem/progenitor\ defects\ in\ TRIM33\ are\ a\ cause\ of\ thyroid\ defects\ defects$ papillary carcinoma (TPC). This antibody is specific to TRIM33. The calculated molecular weight of TRIM33 is 122kDa, but the molecular weight detected is about 140-150 kDa the same as the result from the paper(PMID:25639486;26592194).


Storage

Storage: Store at -80°C. Storage Buffer: PBS Only


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 85364-3-RR (TRIM33 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 85364-3-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using TRIM33 antibody (85364-3-RR, Clone: 242728H2) at dilution of 1:500 and Coralite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) (5A00013-2), CL594-Phalloidin (red). This data was developed using the same antibody clone with 85364-3-PBS in a different storage buffer formulation.



COLO 320 cells were subjected to SDS PAGE followed by western blot with 85364-3-RR (TRIM33 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 85364-3-PBS in a different storage buffer formulation.