## For Research Use Only

## VAMP8 Recombinant antibody

Catalog Number:85400-4-RR



**Basic Information** 

Catalog Number: GenBank Accession Number: **Purification Method:** 85400-4-RR BC001634 Protein A purification

GeneID (NCBI): CloneNo.: 100ul , Concentration: 1000  $\mu g/ml$  by 8673 242830B11

Nanodrop: **UNIPROT ID:** Recommended Dilutions: Source Q9BV40 WB 1:5000-1:50000 Rabbit IHC 1:1000-1:4000 Full Name: vesicle-associated membrane protein F/ICC 1:200-1:800 Isotype:

8 (endobrevin) IgG Immunogen Catalog Number: Calculated MW: AG7903 11 kDa

> Observed MW: 15 kDa

**Applications** 

**Tested Applications:** WB, IHC, IF/ICC, ELISA

Species Specificity: human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: THP-1 cells, HL-60 cells, A431 cells, HeLa cells, rat

kidney tissue, mouse kidney tissue

IHC: human placenta tissue,

IF/ICC: THP-1 cells,

## **Background Information**

VAMP8, also named as endobrevin, is a member of the vesicle-associated membrane protein (VAMP)/synaptobrevin family and the SNARE (soluble NSF-attachment protein receptor) superfamily. Characterized by a common sequence called the SNARE motif, SNARE proteins are involved in membrane fusion and vesicular transport (PMID: 11252968). VAMP8 is involved in autophagy through the direct control of autophagosome membrane fusion with the lysososome membrane. It is required for dense-granule secretion in platelets and plays a role in regulated enzyme secretion in pancreatic acinar cells. VAMP8 is also involved in the abscission of the midbody during cell division, which leads to completely separate daughter cells.

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


## **Selected Validation Data**




Various lysates were subjected to SDS PAGE followed by western blot with 85400-4-RR (VAMP8 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human placenta tissue slide using 85400-4-RR (VAMP8 antibody) at dilution of 1:2000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human placenta tissue slide using 85400-4-RR (VAMP8 antibody) at dilution of 1:2000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed THP-1 cells using VAMP8 antibody (85400-4-RR, Clone: 242830B11) at dilution of 1:400 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2).



Biolayer interferometry (BLL) kinetic assays of 85400-4-RR against Human VAMP8 were performed. The affinity constant is 0.631 nM.