## For Research Use Only

## CDO1 Recombinant monoclonal antibody

Catalog Number:85636-2-RR



**Basic Information** 

Catalog Number: GenBank Accession Number:

85636-2-RR BC024241 GeneID (NCBI): 100ul , Concentration: 1000  $\mu g/ml$  by 1036

Nanodrop: **UNIPROT ID:** Q16878 Rabbit Full Name:

Isotype: cysteine dioxygenase, type I

IgG Calculated MW: Immunogen Catalog Number: 200 aa, 23 kDa AG3294 Observed MW: 23 kDa

**Purification Method:** 

250039D9

Protein A purification CloneNo.:

Recommended Dilutions: WB: 1:1000-1:8000

**Applications** 

**Tested Applications:** 

WB, ELISA

Species Specificity: human, mouse, rat

Positive Controls:

WB: rat liver tissue, mouse liver tissue

## **Background Information**

CDO1(cysteine dioxygenase type 1) is also named as CDO and belongs to the cysteine dioxygenase family. It is an enzyme that adds molecular oxygen to the sulfur of cysteine, converting the thiol to a sulfinic acid known as cysteinesulfinic acid (3-sulfinoalanine) and CDO1 is one of the most highly regulated metabolic enzymes responding to diet(PMID:19011731). The expression of CDO can significantly decrease the level of intracellular cysteine and this reduction is also associated with a decrease in total glutathione levels(PMID:17327371). Cysteine dioxygenase (CDO) plays a critical role in the regulation of cellular cysteine concentration and multiple forms of CDO (23 kDa, 25 kDa, and 68 kDa) have been claimed based upon separation and detection using SDS-PAGE/western blotting(PMID: 14752623).

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3


Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


## **Selected Validation Data**



Various lysates were subjected to SDS PAGE followed by western blot with 85636-2-RR (CDO 1 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.



rat liver tissue were subjected to SDS PAGE followed by western blot with 85636-2-RR (CDO 1 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.



Biolayer interferometry (BLI) kinetic assays of 85636-2-RR against Human CDO1 were performed. The affinity constant is 0.936 nM.