For Research Use Only

OCT1 Recombinant antibody

Catalog Number:85726-1-RR



Purification Method:

Protein A purification

Recommended Dilutions:

WB 1:2000-1:10000

IF/ICC 1:400-1:1600

CloneNo.:

243074D12

Basic Information

Catalog Number: GenBank Accession Number:

85726-1-RR BC003571
Size: GeneID (NCBI):

100ul , Concentration: 1000 µg/ml by 5451
Nanodrop; UNIPROT ID:
Source: P14859
Rabbit Full Name:

Isotype: POU class 2 homeobox 1

IgG Calculated MW:
Immunogen Catalog Number: 76 kDa

Immunogen Catalog Number:76 kDaAG0586Observed MW:

90 kDa

Applications

Tested Applications: Positive Controls:

WB, IF/ICC, ELISA WB: A431 cells, HepG2 cells, HeLa cells, NCCIT cells,

Species Specificity: HEK-293 cells human IF/ICC: HeLa cells,

Background Information

The POU class 2 homeobox 1 (POU2F1), also known as octamer-binding transcription factor-1 (OCT-1), is a ubiquitous nuclear protein that regulates the transcription of target genes by binding to a cis-acting octamer element(5'-ATTTGCAT-3'). It participated in the regulation of housekeeping genes, including histone H2B and snRNAs, for cell differentiation and proliferation [PMID:3462701]. In addition, it functions as a cellular stress sensor that modulates activities of tissue-specific target genes involved in immune and inflammatory responses [PMID:16322220]. It also acts as a repressor for the cytokine-inducible, NF-kB-dependent expression of E-selectin and the vascular cell adhesion molecule during vascular inflammation. Otherwise, it can interact with nuclear hormone receptors, such as the retinoid X receptor, the thyroid hormone receptor and the glucocorticoid receptor, and influence their transcription activity[PMID:17192276].

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 85726-1-RR (OCT1 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using OCT1 antibody (85726-1-RR, Clone: 243074D12) at dilution of 1:800 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-Phalloidin (red).



Biolayer interferometry (BLI) kinetic assays of 85726-1-RR against Human OCT1 were performed. The affinity constant is below 1 pM.