

For Research Use Only

TAP1 Recombinant antibody

Catalog Number: 85766-4-RR



Basic Information

Catalog Number:

85766-4-RR

Size:

100ul, Concentration: 1000 µg/ml by
Nanodrop;

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG1619

GenBank Accession Number:

BC014081

GeneID (NCBI):

6890

UNIPROT ID:

Q03518

Full Name:

transporter 1, ATP-binding cassette,
sub-family B (MDR/TAP)

Calculated MW:

81 kDa

Observed MW:

70-80 kDa

Purification Method:

Protein A purification

CloneNo.:

250037E7

Recommended Dilutions:

WB 1:2000-1:10000

IP 0.5-4.0 µg for 1.0-3.0 mg of total
protein lysate

Applications

Tested Applications:

WB, IP, ELISA

Species Specificity:

human, mouse

Positive Controls:

WB : HeLa cells, HCT 116 cells, THP-1 cells, HepG2
cells, mouse spleen tissue

IP : HepG2 cells,

Background Information

TAP1, also known as ABCB2, PSF1 or RING4, is a member of the ATP-binding cassette (ABC) family of transmembrane transporters and is an essential component of the major histocompatibility complex (MHC) class I antigen-presenting pathway. TAP is involved in the transport of antigens from the cytoplasm to the endoplasmic reticulum for association with MHC class I molecules. It also acts as a molecular scaffold for the final stage of MHC class I folding. Defects in TAP1 are a cause of bare lymphocyte syndrome type 1 (BLS1). Western blot analysis using this antibody detected a major band around 70-80 kDa in HeLa cells.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 85766-4-RR (TAP1 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



THP-1 cells were subjected to SDS PAGE followed by western blot with 85766-4-RR (TAP1 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



HCT 116 cells were subjected to SDS PAGE followed by western blot with 85766-4-RR (TAP1 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



IP result of anti-TAP1 (IP:85766-4-RR, 4ug; Detection:85766-4-RR 1:1000) with HepG2 cells lysate 1800 ug.