

For Research Use Only

Phospho-Myosin Light Chain 2 (Thr18/Ser19) Recombinant antibody

Catalog Number: 85804-1-RR



Basic Information

Catalog Number:

85804-1-RR

Size:

100ul, Concentration: 1000 µg/ml by
Nanodrop;

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

BC016372

GeneID (NCBI):

10627

UNIPROT ID:

P19105

Full Name:

myosin, light chain 12A, regulatory,
non-sarcomeric

Calculated MW:

171 aa, 20 kDa

Observed MW:

19-20 kDa

Purification Method:

Protein A purification

CloneNo.:

250098F12

Recommended Dilutions:

WB 1:500-1:2000

Applications

Tested Applications:

WB, ELISA

Species Specificity:

human, mouse, rat

Positive Controls:

WB: Calyculin A treated NIH/3T3 cells, Calyculin A
treated HEK-293 cells, Calyculin A treated HSC-T6
cells

Background Information

Myosin II is an actin-binding protein composed of MHC (myosin heavy chain), MLCs/MRLCs (myosin regulatory light chains) and ELCs (essential light chains). The Myosin Light Chains are divided into basic MLC (MLC I) and regulatory MLC (MLC II or Myosin Regulatory Light Chain, MRLC). MRLC can be classified into two major types: class I MRLC found in muscles (MLC1, MLC2, MLC3, MYL4, MYL5, MYL6, MYL7, etc), and class II MRLCs present in non-muscle cells (such as MYL9, MYL12A, MYL12B). Phosphorylation of myosin II regulatory light chain (MRLC) is important for cell motility and cytokinesis in nonmuscle cells. Phosphorylation of MRLC at Ser19 (1P-MRLC) enhances myosin II activity, which is further enhanced by MRLC diphosphorylation at Thr18/Ser19 (2P-MRLC) in vitro. PMID: 23951055, PMID: 21126233

This antibody can recognize the phosphorylation of three highly homologous Myosin-regulated light chains: MYL12A, MYL12B and MYL9.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech
Group brand and is not available to purchase from any
other manufacturer.


Selected Validation Data



Calyculin A treated HEK-293 cells were subjected to SDS PAGE followed by western blot with 85804-1-RR (Phospho-Myosin Light Chain 2 (Thr18/Ser19) antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Calyculin A treated NIH/3T3 cells were subjected to SDS PAGE followed by western blot with 85804-1-RR (Phospho-Myosin Light Chain 2 (Thr18/Ser19) antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Calyculin A treated HSC-T6 cells were subjected to SDS PAGE followed by western blot with 85804-1-RR (Phospho-Myosin Light Chain 2 (Thr18/Ser19) antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.