

For Research Use Only

TGFB111 Recombinant antibody, PBS Only

Catalog Number: 85859-5-PBS



Basic Information

Catalog Number: 85859-5-PBS	GenBank Accession Number: BC001830	Purification Method: Protein A purification
Size: 100ug , Concentration: 1 mg/ml by Nanodrop;	GeneID (NCBI): 7041	CloneNo.: 250139E8
Source: Rabbit	UNIPROT ID: O43294	
Isotype: IgG	Full Name: transforming growth factor beta 1 induced transcript 1	
Immunogen Catalog Number: AG0868	Calculated MW: 48 kDa	
	Observed MW: 48-50 kDa	

Applications

Tested Applications:
WB, Indirect ELISA

Species Specificity:
human, rat

Background Information

HIC5 (Hydrogen Peroxide-Inducible Clone 5), also known as TGFB111, ARA55, or TSC-5. It was first identified in 1994 as a gene induced by transforming growth factor- β (TGF- β) or hydrogen peroxide (PMID: 36222304). HIC5 is a member of the Paxillin protein family and functions as a molecular adapter, delivering signals when the cellular adhesion environment changes. Given its role in cancer and fibrotic diseases, HIC5 is a potential therapeutic target. For example, targeting HIC5 or the GR tau2 region could modulate the set of genes regulated by GR, providing opportunities for intervention (PMID: 24591583).

Storage


Storage:
Store at -80°C.

Storage Buffer:
PBS only, pH7.3


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



HeLa cells were subjected to SDS PAGE followed by western blot with 85859-5-RR (TGFβ11 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 85859-5-PBS in a different storage buffer formulation.



SKOV-3 cells were subjected to SDS PAGE followed by western blot with 85859-5-RR (TGFβ11 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 85859-5-PBS in a different storage buffer formulation.



rat colon tissue were subjected to SDS PAGE followed by western blot with 85859-5-RR (TGFβ11 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 85859-5-PBS in a different storage buffer formulation.