For Research Use Only

FABP4 Recombinant antibody

Catalog Number:85864-2-RR



Basic Information

Catalog Number: GenBank Accession Number:

85864-2-RR NM_001442.3

GeneID (NCBI): Size: CloneNo.: 100ul , Concentration: 1000 $\mu g/ml$ by 2167 250188G8

Nanodrop: **UNIPROT ID:** Recommended Dilutions: P15090 WB: 1:5000-1:50000

Rabbit Full Name:

Isotype: fatty acid binding protein 4, adipocyte

IgG Calculated MW:

Immunogen Catalog Number: 15 kDa

EG0880 Observed MW:

15 kDa

Applications

Tested Applications:

WB, ELISA

Species Specificity: human, mouse, rat

Positive Controls:

WB: human heart tissue, human adipose tissue, mouse adipose tissue, rat adipose tissue, mouse brown adipose tissue, mouse heart tissue, mouse small

Purification Method:

Protein A purification

intestine tissue

Background Information

Fatty acid binding protein (FABP) 4 is a member of the FABP family which abundantly expressed, fatty acid carrier proteins. FABPs are capable of binding a variety of hydrophobic molecules such as long-chain fatty acids and are important for their uptake and intracellular trafficking. It was first identified as an adipocyte-specific protein, important for the maintenance of lipid and glucose metabolism. It is also detected in macrophages, where it participates in regulating inflammation and cholesterol trafficking via NFkB and PPAR. In more recent studies, FABP4 has been found in a variety of endothelial cells, where it has been identified as a target of VEGF and a regulator of cell proliferation and possibly angiogenesis. Pathologically, FABP4 has been associated with the development of metabolic syndrome, diabetes and cancer and vulnerability of atherosclerotic plaques. FABP4 has been identified as a novel prognostic factor for both adverse cardiovascular events and breast cancer.

Storage


Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 85864-2-RR (FABP4 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.