## For Research Use Only

## SYNJ2BP Recombinant antibody

Catalog Number:85870-4-RR



**Basic Information** 

Catalog Number: GenBank Accession Number:

85870-4-RR GeneID (NCBI): Size:

100ul , Concentration: 1000  $\mu g/ml$  by 55333 Nanodrop;

**UNIPROT ID:** Source: P57105 Rabbit Full Name:

Isotype synaptojanin 2 binding protein

IgG Calculated MW: Immunogen Catalog Number: 145 aa, 16 kDa AG8130 Observed MW:

16 kDa

**Applications** 

**Tested Applications:** 

WB, IF/ICC, ELISA

Species Specificity:

human, rat

Positive Controls:

WB: A549 cells, rat kidney tissue, SGC-7901 cells, SH-

**Purification Method:** 

CloneNo.:

250205D7

Protein A purification

**Recommended Dilutions:** WB 1:2000-1:10000

IF/ICC 1:250-1:1000

SY5Y cells, human heart tissue

IF/ICC: HeLa cells.

## **Background Information**

Activin, a member of the TGF-beta superfamily, plays important roles in reproductive tissues as a stimulator of follicle-stimulating hormone (FSH) secretion and inhibits the proliferation of breast cancer cells. Activin receptorinteracting protein 2 (ARIP2), also known as synaptojanin-2-binding protein (SYNJ2BP), is located in the outer membrane of mitochondria. SYNJ2BP is widely distributed in various human tissues, and has been recently identified in mouse tissues as a regulatory protein of activin signal transduction by interacting with activin type II and the property of treceptors. SYNJ2BP may be a putative growth-promoting factor involved in breast tumorigenesis and tumor development. A double band of size 15.9 kDa can be detected in Western blot (PMID: 23284606).

Storage

Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol, pH7.3


\*\*\* 20ul sizes contain 0.1%BSA

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**



Rat kidney tissue lysates were subjected to SDS PAGE followed by western blot with 85870-4-RR (ARIP2; SYNJ2BP antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 85870-4-RR (ARIP2; SYNJ2BP antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using ARIP2; SYN12BP antibody (85870-4-RR, Clone: 250205D7) at dilution of 1:500 and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-Phalloidin (red).