For Research Use Only

Cyclin B1 Recombinant antibody, PBS Only




Catalog Number:85895-3-PBS


Basic Information	Catalog Number:	GenBank Accession Number:	Purification Method:
	85895-3-PBS	BC006510	Protein A purification
	Size:	GenelD (NCBI):	CloneNo.:
	100ug, Concentration: 1 mg/ml by	891	250227C12
	Nanodrop; Source: Rabbit Isotype:	UNIPROT ID:	
		P14635	
		Full Name:	
		cyclin B1	
	IgG	Calculated MW:	
	Immunogen Catalog Number: AG29426	48 kDa	
		Observed MW:	
		55-60 kDa	
Applications	Tested Applications: WB, Indirect ELISA Species Specificity: human		
Background Information	Cyclin B1 is a regulatory protein involved in mitosis. The gene product complexes with p34(cdc2) to form the maturation-promoting factor (MPF). Two alternative transcripts have been found, a constitutively expressed transcript and a cell cycle-regulated transcript, that is expressed predominantly during G2/M phase of the cell cycle. The different transcripts result from the use of alternate transcription initiation sites. The antibody is specific to CCNB1. We got a 55-60 kDa band in western blotting maybe due to phosphorylation.		
Storage	Storage: Store at -80°C. Storage Buffer: PBS only, pH7.3		

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com


This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





GE followed by Al antibody) at m temperature ed using the PBS in a Carter of the same antibody clone with 85895-3-PBS in a different storage buffer formulation. Various lysates were subjected to SDS PAGE followed by western blot with 85895-3-RR (CCNB1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 85895-3-PBS in a different storage buffer formulation.



Biolayer interferometry (BLI) kinetic assays of 85895-3-RR against Human Cyclin B1 were performed. The affinity constant is 2.70 nM.

HeLa cells were subjected to SDS PAGE followed by western blot with 85895-3-RR (CCNB1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 85895-3-PBS in a different storage buffer formulation.