For Research Use Only

TIMP1 Recombinant antibody

Catalog Number:85959-1-RR



Basic Information

Catalog Number: GenBank Accession Number:

85959-1-RR NM_001044384.1 Protein A purification

GeneID (NCBI): CloneNo.: 100ul , Concentration: 1000 $\mu g/ml$ by 21857 243010H8

Nanodrop: **UNIPROT ID:** Recommended Dilutions: P12032 WB: 1:5000-1:50000 Rabbit IHC: 1:250-1:1000 Full Name: tissue inhibitor of metalloproteinase IF/ICC: 1:50-1:500

Isotype: IgG

Immunogen Catalog Number: Calculated MW:

23KD EG0630

> Observed MW: 20-23 kDa

Applications

Tested Applications: WB, IHC, IF/ICC, ELISA

Species Specificity:

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: NIH/3T3 cells, C2C12 cells, mouse ovary tissue

Purification Method:

IHC: mouse kidney tissue, IF/ICC: C2C12 cells,

Background Information

TIMP1 is a member of the family of matrix metalloproteinase inhibitors, which contains four members (TIMP1, TIMP2, TIMP3, and TIMP4). Tissue inhibitors of metalloproteinases (TIMPs) are multifaceted molecules that exhibit properties beyond their classical proteinase inhibitory function. TIMP1 has several MMP-independent functions such as modulation of angiogenesis, promotion of cell proliferation, and inhibition of apoptosis. TIMP1 plays important role in cell cycle regulation and cancer progression. Recently, clinical studies have shown that the aberrant $expression \ of \ TIMP1\ is\ associated\ with\ an\ unfavorable\ prognosis\ in\ a\ series\ of\ tumors,\ such\ as\ gastric\ cancer,$ papillary thyroid carcinoma, cutaneous melanoma and breast cancer. In pregnancy, TIMP1 plays a regulatory role in the process of implantation, particularly the cytotrophoblast invasion of the uterine endometrium. In pregnancy, TIMP1 plays a regulatory role in the process of implantation, particularly the cytotrophoblast invasion of the uterine endometrium.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 85959-1-RR (Timp1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed C2C12 cells using TIMP1 antibody (85959-1-RR, Clone: 243010H8) at dilution of 1:200 and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2).



Immunohistochemical analysis of paraffinembedded mouse kidney tissue slide using 85959-1-RR (TIMP1 antibody) at dilution of 1:500 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Biolayer interferometry (BLI) kinetic assays of 85959-1-RR against Mouse TIMP1 were performed. The affinity constant is below 1 pM.