

For Research Use Only

RARG Recombinant antibody

Catalog Number: 86098-1-RR



Basic Information

Catalog Number:

86098-1-RR

Size:

100ul, Concentration: 1000 µg/ml by Nanodrop;

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG1991

GenBank Accession Number:

BC019098

GeneID (NCBI):

5916

UNIPROT ID:

P13631

Full Name:

retinoic acid receptor, gamma

Calculated MW:

454 aa, 50 kDa

Observed MW:

50-60kDa

Purification Method:

Protein A purification

CloneNo.:

250399C1

Recommended Dilutions:

WB: 1:500-1:2000

Applications

Tested Applications:

WB, ELISA

Species Specificity:

human

Positive Controls:

WB : MCF-7 cells, HeLa cells, NCCIT cells, T-47D cells, A431 cells

Background Information

Retinoic acid receptors (RARs) are nuclear hormone receptors that act as ligand-dependent transcriptional regulators. In their liganded state, RARs activate transcription, whereas in their nonliganded form, they repress transcription of their target genes. RARs have numerous target genes, which have retinoic response elements in their promoter regions (PMID:17574023). RARG is a member of the retinoid acid receptor (RAR) family, along with RARA, which is known to be involved PML/RARA fusion of acute promyelocytic leukemia, and RARB. and rearrangement that conferred to leukemic cells acute promyelocytic leukemia-like morphologic and immunophenotypic features (PMID:20935257).

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 86098-1-RR (RARG antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.