## For Research Use Only

## MGMT Recombinant antibody

Catalog Number:86115-5-RR



**Basic Information** 

Catalog Number: GenBank Accession Number:

86115-5-RR BC000824 Protein A purification GeneID (NCBI): CloneNo.: 100ul , Concentration: 1000  $\mu g/ml$  by 4255 243130G10

Nanodrop: **UNIPROT ID:** P16455 Rabbit Full Name:

Isotype: O-6-methylguanine-DNA methyltransferase IgG Immunogen Catalog Number: Calculated MW: AG9996 22 kDa

> Observed MW: 22 kDa

**Applications** 

**Tested Applications:** WB, IHC, IF/ICC, ELISA

Species Specificity:

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

WB: Jurkat cells, MCF-7 cells, Raji cells, HeLa cells,

**Purification Method:** 

Recommended Dilutions:

WB: 1:2000-1:10000

IF/ICC: 1:400-1:1600

IHC: 1:500-1:2000

HepG2 cells, LNCaP cells

IHC: human lung cancer tissue,

IF/ICC: MCF-7 cells.

## **Background Information**

MGMT is the primary vehicle for cellular removal of alkyl lesions from the O-6 position of guanine and the O-4 position of thymine. While key to the maintenance of genomic integrity, MGMT also removes damage induced by alkylating chemotherapies, inhibiting the efficacy of cancer treatment (PMID:23065697). MGMT is the primary mechanism for the removal of alkylation damage from the O-6 position of guanine (PMID: 17482892). The O-6 position of guanine is one of several positions in DNA bases to which alkyl groups are attached in SN1 alkylation reactions, and this repair has been well-characterized in mammalian cells and via MGMT homologs in bacteria and Archaea (PMID: 10767620).

Storage


Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


## **Selected Validation Data**



Various lysates were subjected to SDS PAGE followed by western blot with 89933-4-RR (MGMT antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.

Immunofluorescent analysis of (4% PFA) fixed MCF-7 cells using MGMT antibody (89933-4-RR, Clone: 243130G10 ) at dilution of 1:800 and CoraLite@488-Conjugated Goat Anti-Rabbit  $\lg G(H+L)$  (SA00013-2), CL594-Phalloidin (red).

Immunohistochemical analysis of paraffinembedded human lung cancer tissue slide using 86115-5-RR (MGMT antibody) at dilution of 1:1000 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human lung cancer tissue slide using 86115-5-RR (MGMT antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).